

GROTHENDIECK GROUPS AND A CATEGORIFICATION OF ADDITIVE INVARIANTS

JÖRG SCHÜRMANN AND SHOJI YOKURA

ABSTRACT. A topologically-invariant and additive homology class is mostly not a natural transformation as it is. In this paper we discuss turning such a homology class into a natural transformation; i.e., a “categorification” of it. In a general categorical set-up we introduce a *generalized relative Grothendieck group* from a cospan of functors of categories and also consider a *categorification of additive invariants on objects*. As an example, we obtain a general theory of characteristic homology classes of singular varieties.

1. INTRODUCTION

Characteristic classes are important invariants in modern geometry and topology to investigate other invariants. Classically characteristic classes with values in cohomology theory were considered for real or complex vector bundles. The main feature of them is that they are formulated as natural transformations from the contravariant functor of vector bundles to the cohomology theory. When it comes to the case of possibly singular varieties, characteristic classes are considered in homology theory, instead of cohomology theory and still formulated as *natural transformations from a covariant functor $\mathcal{F}$ to a (suitable) homology theory H_* , satisfying the normalization condition that for a smooth variety X the value of a distinguished element Δ_X of $\mathcal{F}(X)$ is equal to the Poincaré dual of the corresponding characteristic cohomology class of the tangent bundle*:

$$\tau_{cl} : \mathcal{F}(-) \rightarrow H_*(-) \quad \text{such that for } X \text{ smooth} \quad \tau_{cl}(\Delta_X) = cl(TX) \cap [X].$$

Most important and well-studied theories of characteristic homology classes of singular varieties are the following ones, where we consider here for simplicity only the category of complex algebraic varieties and functoriality is required only for proper morphisms. Below, in (1) and (2) the homology theory $H_*(X)$ is either Chow groups $CH_*(X)$ or the even-degree Borel–Moore homology groups $H_{2*}^{BM}(X)$, whereas in (3) X is assumed to be compact and thus the homology theory $H_*(X)$ is the usual even-degree homology group $H_{2*}(X)$:

- (1) MacPherson’s Chern class [Mac1] $c_*^{Mac} : F(-) \rightarrow H_*(-)$ is the unique natural transformation from the covariant functor F of constructible functions to the homology H_* , satisfying the normalization condition that for a smooth variety X the value of the characteristic function is the Poincaré dual of the total Chern class of the tangent bundle:

$$c_*^{Mac}(\mathbb{1}_X) = c(TX) \cap [X].$$

- (2) Baum–Fulton–MacPherson’s Todd class [BFM] $td_*^{BFM} : G_0(X) \rightarrow H_*(X) \otimes \mathbb{Q}$ is the unique natural transformation from the covariant functor G_0 of Grothendieck groups of coherent sheaves to the rationalized homology $H_* \otimes \mathbb{Q}$, satisfying the normalization condition that for a smooth variety X the value of (the class of) the structure sheaf is the Poincaré dual of the total Todd class of the tangent bundle:

$$td_*^{BFM}([\mathcal{O}_X]) = td(TX) \cap [X].$$

(1) Jörg Schürmann: supported by the SFB 878 “groups, geometry and actions”.

(2) Shoji Yokura: partially supported by Grant-in-Aid for Scientific Research (No. 21540088), the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan

Keywords: homology class, characteristic homology class, categorification, geometric homology, bordism, Steenrod’s problem.

Mathematics Subject Classification 2000: 55Nxx, 14C17, 14C40, 18A99.

- (3) Goresky–MacPherson’s homology L -class [GM], which is extended as a natural transformation by Cappell–Shaneson [CS] (also see [Yo2]), $L_*^{CS} : \Omega(X) \rightarrow H_*(X) \otimes \mathbb{Q}$ is a natural transformation from the covariant functor Ω of cobordism groups of self-dual constructible sheaf complexes to the rationalized homology theory $H_* \otimes \mathbb{Q}$, satisfying the normalization condition that for a smooth variety X the value of the (class of the) shifted constant sheaf is the Poincaré dual of the total Hirzebruch–Thom’s L -class of the tangent bundle:

$$L_*^{CS}([\mathbb{Q}_X[\dim X]]) = L(TX) \cap [X].$$

Here X is assumed to be compact.

Recently these three theories c_*^{Mac} , td_*^{BFM} and L_*^{CS} are “unified” by the motivic Hirzebruch class [BSY] (see also [SY], [Sch] and [Yo4]) $T_{y*} : K_0(\mathcal{V}_{\mathbb{C}}/X) \rightarrow H_*(X) \otimes \mathbb{Q}[y]$, which is the unique natural transformation such that for a smooth variety X the value of the isomorphism class of the identity $X \xrightarrow{id_X} X$ is the Poincaré dual of the generalized Hirzebruch–Todd class $T_y(TX)$ of the tangent bundle:

$$T_{y*}([X \xrightarrow{id_X} X]) = T_y(TX) \cap [X].$$

Here $K_0(\mathcal{V}_{\mathbb{C}}/X)$ is the relative Grothendieck group of the category $\mathcal{V}_{\mathbb{C}}$ of complex algebraic varieties, i.e., the free abelian group generated by the isomorphism classes $[V \xrightarrow{h} X]$ of morphism $h \in \text{hom}_{\mathcal{V}_{\mathbb{C}}}(V, X)$ modulo the relations

- $[V_1 \xrightarrow{h_1} X] + [V_2 \xrightarrow{h_2} X] = [V_1 \sqcup V_2 \xrightarrow{h_1+h_2} X]$, with $\sqcup$ the disjoint union, and
- $[V \xrightarrow{h} X] = [V \setminus W \xrightarrow{h_V \setminus W} X] + [W \xrightarrow{h_W} X]$ for $W \subset V$ a closed subvariety of V .

The generalized Hirzebruch–Todd class $T_y(E)$ of the complex vector bundle E (see [Hir, HBJ]) is defined to be

$$T_y(E) := \prod_{i=1}^{\text{rank } E} \left(\frac{\alpha_i(1+y)}{1 - e^{-\alpha_i(1+y)}} - \alpha_i y \right) \in H^*(X) \otimes \mathbb{Q}[y],$$

where α_i is the Chern root of E , i.e., $c(E) = \prod_{i=1}^{\text{rank } E} (1 + \alpha_i)$. Note that

- (1) $T_{-1}(E) = c(E)$ is the Chern class,
- (2) $T_0(E) = td(E)$ is the Todd class and
- (3) $T_1(E) = L(E)$ is the Thom–Hirzebruch L -class.

The “unification” means the existence of the following commutative diagrams of natural transformations:

$$\begin{array}{ccccc} K_0(\mathcal{V}_{\mathbb{C}}/X) & & K_0(\mathcal{V}_{\mathbb{C}}/X) & & K_0(\mathcal{V}_{\mathbb{C}}/X) \\ \downarrow T_{-1*} & \searrow \text{const} & \downarrow T_{0*} & \searrow \text{coh} & \downarrow T_{1*} \\ & F(X) & & G_0(X) & \Omega(X) \\ & \swarrow c_*^{Mac} \otimes \mathbb{Q} & & \swarrow td_*^{BFM} & \swarrow L_*^{CS} \\ H_*(X) \otimes \mathbb{Q} & & H_*(X) \otimes \mathbb{Q} & & H_*(X) \otimes \mathbb{Q} \end{array}$$

Here $\text{const} : K_0(\mathcal{V}_{\mathbb{C}}/X) \rightarrow F(X)$ is defined by $\text{const}([V \xrightarrow{h} X]) := h_* \mathbb{1}_V$. The other two comparison transformations are characterized by $\text{coh}([V \xrightarrow{h} X]) = h_*([O_V])$ and $\text{sd}([V \xrightarrow{h} X]) = h_*([\mathbb{Q}_V[\dim V]])$ for V smooth and h proper. $T_{y*}(X) := T_{y*}([X \xrightarrow{id_X} X])$ is called the *motivic Hirzebruch class of X* . For more details see [BSY].

Besides these characteristic classes formulated as natural transformations, there are several important homology classes which are usually not formulated or not captured as such natural transformations; for example: Chern–Mather class $c_*^M(X)$ ([Mac1]), Segre–Mather class $s_*^M(X)$ ([Yo1]), Fulton’s canonical Chern class $c_*^F(X)$ ([Fu]), Fulton–Johnson’s Chern class $c_*^{FJ}(X)$ ([FJ, Fu]), Milnor class $\mathcal{M}(X)$ (e.g., see [Al], [BLSS], [PP], [Yo3], etc.), which is (up to sign) the difference of the MacPherson’s Chern class and the Fulton–Johnson’s Chern class, etc.

In [Yo5] we captured Fulton–Johnson’s Chern class and the Milnor class $\mathcal{M}(X)$ as natural transformations, with the latter one as a special case of the Hirzebruch–Milnor class (also see [CMSS]), using the motivic Hirzebruch class. In this paper we generalize the approach and results of [Yo5] to a more general abstract categorical context (cf. [Yo6]).

Let $\mathcal{B}$ be a category with a *coproduct* $\sqcup$. Here we assume that a fixed coproduct $X \sqcup Y$ of any two objects $X, Y \in \text{ob}(\mathcal{B})$ has been chosen, as well as an initial object $\emptyset$ in $\mathcal{B}$, so that we can view $(\mathcal{B}, \sqcup)$ as a symmetric monoidal category with unit $\emptyset$. Examples are the category $\mathcal{TOP}$ (resp., $\mathcal{TOP}_{lc}$) of (locally compact) topological spaces, the category $\mathcal{C}^\infty$ of $\mathcal{C}^\infty$ -manifolds, or the category $\mathcal{V}_k$ of algebraic varieties (i.e., reduced separated schemes of finite type) over a base field k , with the coproduct $\sqcup$ given by the usual disjoint union. Moreover, for $\mathcal{B} = \mathcal{TOP}_{lc}, \mathcal{C}^\infty$ or $\mathcal{V}_k$, we often consider them as categories only with respect to proper morphisms. Let $\mathcal{AB}$ be the category of abelian groups with the coproduct $\sqcup$ given by the direct sum $\oplus$. Then a covariant functor $\mathcal{H} : \mathcal{B} \rightarrow \mathcal{AB}$ is called *additive* (cf. [LM]), if it preserves the coproduct structure $\sqcup$, i.e., if we get the following isomorphism:

$$(i_X)_* \oplus (i_Y)_* : \mathcal{H}(X) \oplus \mathcal{H}(Y) \xrightarrow{\cong} \mathcal{H}(X \sqcup Y)$$

where $i_X : X \rightarrow X \sqcup Y$ and $i_Y : Y \rightarrow X \sqcup Y$ are the canonical injections. In particular, $\mathcal{H}(\emptyset) = \{0\}$. Examples of such an additive functor $\mathcal{H}$ are a generalized homology theory H_* for $\mathcal{B} = \mathcal{TOP}$, the Borel–Moore homology theory H_*^{BM} for $\mathcal{B} = \mathcal{TOP}_{lc}$, the functor G_0 of Grothendieck groups of coherent sheaves or the Chow groups CH_* for $\mathcal{B} = \mathcal{V}_k$, as well as the functor F of constructible functions or Ω of cobordism groups of self-dual constructible sheaf complexes for $\mathcal{B} = \mathcal{V}_{\mathbb{C}}$.

Let $\mathcal{C}$ be a category of “(topological) spaces with some additional structures *stable* under $\sqcup$ ”, such as the category of (complex) algebraic varieties, the category of (compact) topological spaces, or the category of oriented smooth (i.e., $\mathcal{C}^\infty$ -) manifolds, etc. More abstractly, $\mathcal{C}$ is a *symmetric monoidal category* $(\mathcal{C}, \sqcup)$ together with a strict monoidal forgetful functor $\mathfrak{f} : (\mathcal{C}, \sqcup) \rightarrow (\mathcal{B}, \sqcup)$. Then an *additive invariant* on objects in $\text{ob}(\mathcal{C})$ with values in the additive functor $\mathcal{H}$ is given by an element $\alpha(X) \in \mathcal{H}(\mathfrak{f}(X))$ for all $X \in \text{ob}(\mathcal{C})$ satisfying

$$\alpha(X \sqcup Y) = (i_{\mathfrak{f}(X)})_* \alpha(X) + (i_{\mathfrak{f}(Y)})_* \alpha(Y),$$

where $i_{\mathfrak{f}(X)}$ and $i_{\mathfrak{f}(Y)}$ are the corresponding canonical inclusions. Note that the sum or difference $\alpha \pm \beta$ of two additive invariants with values in $\mathcal{H}$ is again such an invariant.

Example 1.1. Let $\mathcal{B} = \mathcal{V}_{\mathbb{C}}$ be the category of complex algebraic varieties and $\mathcal{C}$ be a full subcategory of $\mathcal{B}$ which is stable under disjoint union, isomorphisms and contains the initial object $\emptyset$ and the final object pt (e.g., an *admissible subcategory* in the sense of Levine–Morel [LM]), such as the category $\mathcal{V}_{\mathbb{C}}^{sm}$ of smooth varieties, $\mathcal{V}_{\mathbb{C}}^{lpd}$ of locally pure dimensional varieties, $\mathcal{V}_{\mathbb{C}}^{emb}$ of varieties embeddable into smooth varieties, or $\mathcal{V}_{\mathbb{C}}^{lci}$ of local complete intersections. Then we have the following additive invariants, some of which are already mentioned above:

- (1) characteristic homology classes $cl_*(X) := cl(TX) \cap [X] \in H_*(X) \otimes R$ for $X \in \text{ob}(\mathcal{V}_{\mathbb{C}}^{sm})$ with $[X]$ the fundamental class and TX the tangent bundle, or the corresponding virtual classes $cl_*(X)$ for $X \in \text{ob}(\mathcal{V}_{\mathbb{C}}^{lci})$, with TX the virtual tangent bundle and $cl(TX)$ a characteristic cohomology class.
- (2) Fulton’s canonical Chern class or Fulton–Johnson Chern class $c_*^F(X)$, $c_*^{FJ}(X)$ for $X \in \text{ob}(\mathcal{V}_{\mathbb{C}}^{emb})$.
- (3) Mather-type characteristic homology classes $cl_*^{Ma}(X)$, such as Chern–Mather class and Segre–Mather class, or the local Euler obstruction $Eu_X \in F(X)$ as a constructible function for $X \in \text{ob}(\mathcal{V}_{\mathbb{C}}^{lpd})$.
- (4) the (class of the) self-dual Intersection Homology complex $[IC_X] \in \Omega(X)$ and for X compact its Goresky–MacPherson’s L -homology class $L^{GM}(X) \in H_*(X)$ for $X \in \text{ob}(\mathcal{V}_{\mathbb{C}}^{lpd})$.

A categorification of the additive invariant α with values in the additive functor $\mathcal{H}$ is meant to be an associated natural transformation

$$\tau_\alpha : \mathcal{H}_\alpha \rightarrow \mathcal{H} \circ \mathfrak{f}$$

from a covariant functor $\mathcal{H}_\alpha$ on the category $\mathcal{C}$, such that

$$\tau_\alpha(\delta_X) = \alpha(X)$$

for some distinguished element $\delta_X \in \mathcal{H}_\alpha(X)$ and all $X \in \text{ob}(\mathcal{C})$.

To construct such a covariant functor $\mathcal{H}_\alpha$, we introduce *generalized relative Grothendieck groups*, using comma categories in a more abstract categorical context. The construction of such a covariant functor is hinted by the definition of the relative Grothendieck group $K_0(\mathcal{V}_\mathbb{C}/X)$ and more clearly by the description of the oriented bordism group $\Omega_*^{SO}(X)$.

The oriented bordism group $\Omega_m^{SO}(X)$ of a topological space X is defined to be the free abelian group generated by the isomorphism classes $[M \xrightarrow{h} X]$ of continuous maps $M \xrightarrow{h} X$ from closed oriented smooth manifolds M of dimension m to the given topological space X , modulo the following relations

- (1) $[M \xrightarrow{h} X] + [M' \xrightarrow{h'} X] = [M \sqcup M' \xrightarrow{h+h'} X]$,
- (2) if $M \xrightarrow{h} X$ and $M' \xrightarrow{h'} X$ are bordant, then $[M \xrightarrow{h} X] = [M' \xrightarrow{h'} X]$.

Note that if $M \xrightarrow{h} X$ and $M' \xrightarrow{h'} X$ are isomorphic to each other, i.e., there exists an orientation preserving diffeomorphism $\phi : M \cong M'$ such that the following diagram commutes as topological spaces:

$$\begin{array}{ccc} M & \xrightarrow{\phi} & M' \\ & \searrow h & \swarrow h' \\ & X, & \end{array}$$

then $M \xrightarrow{h} X$ and $M' \xrightarrow{h'} X$ are clearly bordant to each other. Hence we can say that the bordism group $\Omega_m^{SO}(X)$ is defined to be the free abelian group generated by isomorphism classes of continuous maps $M \xrightarrow{h} X$ from closed oriented smooth manifolds M of dimension m to the given topological space X , modulo the relations (1) and (2) above. The bordism group $\Omega_m^{SO}(X)$ can also be described slightly differently as follows: the set of the bordism classes $[M \xrightarrow{h} X]$ can be turned into a commutative semi-group or monoid $\mathcal{M}(X)$ by the relation (1) above, with $0 = [\emptyset \rightarrow X]$. Then the Grothendieck group or the group completion of the monoid $\mathcal{M}(X)$ is nothing but the above bordism group $\Omega_m^{SO}(X)$.

In the definition of the bordism group two categories are involved:

- the category $\mathcal{C}_{co}^\infty$ of closed oriented $\mathcal{C}^\infty$ -manifolds, i.e., compact oriented $\mathcal{C}^\infty$ -manifolds without boundary,
- the category $\mathcal{TOP}$ of topological spaces.

It should be emphasized that *even though we consider such a finer category $\mathcal{C}_{co}^\infty$ for a source space M the map $h : M \rightarrow X$ of course has to be considered in the crude category $\mathcal{TOP}$.*

The bordism group Ω_*^{SO} is a generalized homology theory, in particular Ω_*^{SO} is a covariant functor

$$\Omega_*^{SO} : \mathcal{TOP} \rightarrow \mathcal{AB},$$

where $\mathcal{AB}$ is the category of abelian groups.

Clearly we can consider this covariant functor on a different category finer than the category $\mathcal{TOP}$ of topological spaces, e.g., the category $\mathcal{V}_\mathbb{C}$ of complex algebraic varieties. Namely we consider continuous maps $h : M \rightarrow V$ from closed oriented manifolds M to a complex algebraic variety V . Then we still get a covariant functor

$$\Omega_*^{SO} : \mathcal{V}_\mathbb{C} \rightarrow \mathcal{AB}.$$

In this set-up three different categories $\mathcal{C}_{co}^\infty$, $\mathcal{TOP}$ and $\mathcal{V}_{\mathbb{C}}$ are involved. More precisely, we have the following forgetful functors $f_s : \mathcal{C}_{co}^\infty \rightarrow \mathcal{TOP}$ and $f_t : \mathcal{V}_{\mathbb{C}} \rightarrow \mathcal{TOP}$ (here “s” and “t” mean “source object” and “target object”):

$$\mathcal{C}_{co}^\infty \xrightarrow{f_s} \mathcal{TOP} \xleftarrow{f_t} \mathcal{V}_{\mathbb{C}}.$$

And the commutative triangle

$$\begin{array}{ccc} M & \xrightarrow{\phi} & M' \\ & \searrow h & \swarrow h' \\ & V & \end{array}$$

really means

$$\begin{array}{ccc} f_s(M) & \xrightarrow{f_s(\phi)} & f_s(M') \\ & \searrow h & \swarrow h' \\ & f_t(V) & \end{array}$$

Abstracting this situation, we deal in this paper with a more general situation of a cospan of categories

$$\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t,$$

where $\mathcal{B}$ is a category with (chosen) coproducts $\sqcup$, $(\mathcal{C}_s, \sqcup)$ is a *symmetric monoidal category* with $S : \mathcal{C}_s \rightarrow \mathcal{B}$ a strict monoidal functor respecting the units $\emptyset$, and $T : \mathcal{C}_t \rightarrow \mathcal{B}$ just a functor.

Two triples $(V, X, h), (V', X, h')$, where $V, V' \in ob(\mathcal{C}_s)$, $X \in ob(\mathcal{C}_t)$, $h \in hom_{\mathcal{B}}(S(V), T(X))$ and $h' \in hom_{\mathcal{B}}(S(V'), T(X))$, are called *isomorphic*, if there exists an isomorphism $\phi : V \xrightarrow{\cong} V' \in hom_{\mathcal{C}_s}(V, V')$ such that the following diagram commutes, just like as above:

$$\begin{array}{ccc} S(V) & \xrightarrow{S(\phi)} & S(V') \\ & \searrow h & \swarrow h' \\ & T(X) & \end{array}$$

Then the isomorphism classes $[(V, X, h)]$ of triples (V, X, h) can be turned into a monoid $\mathcal{M}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$ by

$$[(V, X, h)] \sqcup [(V', X, h')] := [(V \sqcup V', X, h + h')],$$

with unit $0 = [(\emptyset, X, h)]$. The associated Grothendieck group is denoted by $K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$ (respectively $K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/X)$ in case $\mathcal{C}_t = \mathcal{B}$ and $T = id_{\mathcal{C}_t}$) and called the *canonical generalized (S, T) -relative Grothendieck group*. It is a covariant functor with respect to X by composing h with $T(f)$ for $f \in hom_{\mathcal{C}_t}(X, X')$.

Remark 1.2. In the following two cases, a *distinguished element* $\delta_X \in K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$ is available:

- (i) $\delta_{S(V)} := [(V, S(V), id_{S(V)})]$ for $X = S(V)$ in the case when $T = id_{\mathcal{C}_t} : \mathcal{C}_t \rightarrow \mathcal{C}_t = \mathcal{B}$ is the identity functor.
- (ii) $\delta_V := [(V, V, id_{S(V)})]$ for $X = V$ in the case when $\mathcal{C}_t = \mathcal{C}_s$ and $T = S$.

When it comes to a categorification of an additive invariant α with values in an additive covariant functor $\mathcal{H} : \mathcal{B} \rightarrow \mathcal{AB}$, we sometimes need to consider a stronger notion of an isomorphism of triples (V, X, h) (e.g. in the context with V an “oriented space”): Let $(V, X, h), (V', X, h')$ be two triples as above. Then they are called *α -isomorphic* if the following holds: they are isomorphic as above by an isomorphism $\phi : V \xrightarrow{\cong} V' \in hom_{\mathcal{C}_s}(V, V')$ such that

$$(S(\phi))_* \alpha(V) = \alpha(V').$$

If this last equality holds for any isomorphism $\phi : V \xrightarrow{\cong} V' \in \text{hom}_{\mathcal{C}_s}(V, V')$, then α is also called an *isomorphism invariant*. So in this case there is no difference between α -isomorphism classes and isomorphism classes for such triples (V, X, h) . The α -isomorphism class of (V, X, h) is denoted by $[(V, X, h)]_\alpha$ and these can be turned as before into a monoid $\mathcal{M}_\alpha(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$, since α is additive. The associated generalized (S, T) -relative Grothendieck group is denoted by $K_\alpha(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$, respectively $K_\alpha(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/X)$ in case $\mathcal{C}_t = B$ and $T = \text{id}_{\mathcal{C}_t}$. If it is clear that we consider this group $K_\alpha(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$ from the context, we sometimes omit the suffix α from the notation, e.g. if α is an isomorphism invariant. Similarly, if we consider $\mathcal{B} = \mathcal{TOP}_{lc}, \mathcal{C}^\infty$ or $\mathcal{V}_k$ only as categories with respect to proper morphisms, then this is indicated by the notation $K_\alpha^{\text{prop}}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$, $K_\alpha^{\text{prop}}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/X)$ etc.

Theorem 1.3. (*A “categorification” of an additive invariant*) Let $\mathcal{H} : \mathcal{B} \rightarrow \mathcal{AB}$ be an additive functor on $\mathcal{B}$ with $T' = \mathcal{H} \circ T$. Then an additive invariant α on $\text{ob}(\mathcal{C}_s)$ with values in $\mathcal{H}$ induces a natural transformation on $\mathcal{C}_t$:

$$\tau_\alpha : K_\alpha(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(-)) \rightarrow T'(-); \tau_\alpha([(V, X, h)]_\alpha) := h_*(\alpha(V)).$$

Assume T is full, and consider one of the cases (i) or (ii) as in Remark 1.2. Then τ_α is the unique natural transformation satisfying

$$\tau_\alpha([(V, S(V), \text{id}_{S(V)})]_\alpha) = \alpha(V) \text{ or } \tau_\alpha([(V, V, \text{id}_{S(V)})]_\alpha) = \alpha(V).$$

Let us illustrate this result by some examples.

Example 1.4. Let $\mathcal{TOP}$ be the category of topological spaces and let $\mathcal{C}_{co}^\infty$ be the category of closed oriented C^∞ -manifolds, whose morphisms are just differentiable maps. Consider the following cospan of these categories:

$$\mathcal{C}_{co}^\infty \xrightarrow{\mathfrak{f}} \mathcal{TOP} \xleftarrow{\text{id}_{\mathcal{TOP}}} \mathcal{TOP},$$

where $\mathfrak{f} : \mathcal{C}_{co}^\infty \rightarrow \mathcal{TOP}$ is the forgetful functor. Then the fundamental class $[M] \in H_*(M)$ for $M \in \text{ob}(\mathcal{C}_{co}^\infty)$ is an additive invariant with values in the usual homology H_* . More generally, let $\text{cl}(E) \in H^*(-; R)$ be a contravariant functorial characteristic class of (isomorphism classes of) oriented vector bundles E . Then also

$$\alpha(M) := \text{cl}(TM) \cap [M] \in H_*(M; R)$$

is additive, where we recover the fundamental class for cl the unit $1 \in H^*(-)$. Assume cl is normalized in the sense that $1 = \text{cl}^0(E) \in H^0(-; R)$, so that a diffeomorphism $\phi : M \xrightarrow{\cong} M'$ of such oriented manifolds is an α -isomorphism if and only if it is orientation preserving. Then there exists a unique natural transformation

$$\tau_{\text{cl}} : K_\alpha(\mathcal{C}_{co}^\infty \xrightarrow{\mathfrak{f}} \mathcal{TOP}/-) \rightarrow H_*(-; R)$$

such that for a closed oriented C^∞ -manifold M

$$\tau_{\text{cl}}([(M, \mathfrak{f}(M), \text{id}_{\mathfrak{f}(M)})]_\alpha) = \text{cl}(TM) \cap [M].$$

Let $\Omega_*^{SO}(X)$ be the oriented bordism group of a topological space X . Then we have

$$\Omega_*^{SO}(X) \cong K_\alpha(\mathcal{C}_{co}^\infty \xrightarrow{\mathfrak{f}} \mathcal{TOP}/X) / \sim,$$

with $\sim$ the bordism relation. And τ_{cl} factorizes over $\Omega_*^{SO}(X)$ in the case when cl is stable in the sense that $\text{cl}(E \oplus \mathbb{R}) = \text{cl}(E)$ for $\mathbb{R}$ the trivial real line bundle.

Example 1.5. Let $\mathcal{V}_{\mathbb{C}}$ be the category of complex algebraic varieties, with $\mathcal{C}$ the full subcategory $\mathcal{V}_{\mathbb{C}}^{\text{sm}}, \mathcal{V}_{\mathbb{C}}^{\text{lpd}}, \mathcal{V}_{\mathbb{C}}^{\text{emb}}$ or $\mathcal{V}_{\mathbb{C}}^{\text{lci}}$ of smooth, locally pure dimensional, embeddable or local complete intersection varieties, with

$$\alpha(V) := \text{cl}_*(V) \in H_*(V) \otimes R$$

for $V \in \text{ob}(\mathcal{C})$ a corresponding additive characteristic homology class as in Example 1.1. Here the homology theory $H_*(X)$ is either Chow groups $CH_*(X)$ or the even-degree Borel–Moore homology groups $H_{2*}^{BM}(X)$.

Then α is also an isomorphism invariant. Consider the following cospan of categories, where we consider only proper morphisms:

$$\mathcal{C} \xrightarrow{f} \mathcal{V}_{\mathbb{C}} \xleftarrow{id_{\mathcal{V}_{\mathbb{C}}}} \mathcal{V}_{\mathbb{C}},$$

with $f : \mathcal{C} \rightarrow \mathcal{V}_{\mathbb{C}}$ being the inclusion functor. Then there exists a unique natural transformation

$$\tau_{cl} : K^{prop}(\mathcal{C} \xrightarrow{f} \mathcal{V}_{\mathbb{C}}/-) \rightarrow H_*(-) \otimes R,$$

such that for all $V \in ob(\mathcal{C})$

$$\tau_{cl}([(V, f(V), id_{f(V)})]) = cl_*(V).$$

Moreover, there is a tautological surjective natural transformation

$$K^{prop}(\mathcal{C} \xrightarrow{f} \mathcal{V}_{\mathbb{C}}/-) \rightarrow K_0(\mathcal{V}_{\mathbb{C}}/-)$$

to the relative Grothendieck group $K_0(\mathcal{V}_{\mathbb{C}}/-)$ of complex algebraic varieties. And for $\mathcal{C} = \mathcal{V}_{\mathbb{C}}^{sm}$ and

$$cl_*(V) = T_y(TV) \cap [V] = T_{y*}(V) \quad \text{or} \quad cl_*(V) = c(TV) \cap [V] = c_*(V),$$

i.e., the Hirzebruch homology class or MacPherson's Chern class, the transformations $\tau_{T_{y*}}$ and τ_{c_*} factorize over $K_0(\mathcal{V}_{\mathbb{C}}/-)$ by [BSY].

In §2 we explain the general categorical background and prove Theorem 1.3, whereas in §§3-4 we apply it to many different geometric situations, e.g., for obtaining Riemann–Roch type theorems.

2. GENERALIZED RELATIVE GROTHENDIECK GROUPS

Definition 2.1. Let $\mathcal{C}$ be a symmetric monoidal category equipped with monoidal structure $\oplus$ and the unit object $\emptyset$. The Grothendieck group $K(\mathcal{C})$ is defined to be the free abelian group generated by the isomorphism classes $[X]$ of objects $X \in ob(\mathcal{C})$ modulo the relations

$$[X] + [Y] = [X \oplus Y], \quad , \text{ with } 0 = [\emptyset].$$

A functor $\Phi : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ of two symmetric monoidal categories is a functor which preserves $\oplus$ in the relaxed sense that there are natural transformations:

$$\Phi(A) \oplus_{\mathcal{C}_2} \Phi(B) \rightarrow \Phi(A \oplus_{\mathcal{C}_1} B),$$

In some usage it requires an isomorphism

$$\Phi(A) \oplus_{\mathcal{C}_2} \Phi(B) \cong \Phi(A \oplus_{\mathcal{C}_1} B),$$

in which case it is called a *strong* monoidal functor. Here we also assume that Φ respects the units.

Example 2.2. Let $\mathcal{B}$ be a category with a (chosen) *coproduct* $\sqcup$ and $\mathcal{H} : \mathcal{B} \rightarrow \mathcal{AB}$ be an additive covariant functor. Then we have

$$\mathcal{H}_*(X \sqcup Y) \cong \mathcal{H}_*(X) \oplus \mathcal{H}_*(Y) \quad , \text{ with } \mathcal{H}_*(\emptyset) = \{0\}.$$

Examples of such an additive functor $\mathcal{H}$ are a generalized homology theory H_* for $\mathcal{B} = \mathcal{TOP}$, the Borel–Moore homology theory H_*^{BM} for $\mathcal{B} = \mathcal{TOP}_{lc}$, the functor G_0 of Grothendieck groups of coherent sheaves or the Chow groups CH_* for $\mathcal{B} = \mathcal{V}_k$, as well as the functor F of constructible functions or Ω of cobordism groups of self-dual constructible sheaf complexes for $\mathcal{B} = \mathcal{V}_{\mathbb{C}}$.

Lemma 2.3. Let $\Phi : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ be a strong monoidal functor of two symmetric monoidal categories. Then the map

$$\Phi_* : K(\mathcal{C}_1) \rightarrow K(\mathcal{C}_2), \quad \Phi_*([X]) := [\Phi(X)]$$

is a well-defined group homomorphism. Namely, the Grothendieck group K is a covariant functor from the category of such categories and functors to the category of abelian groups.

Now we recall the notion of *comma category* and *fiber category* (e.g., see [MacL]):

Definition 2.4. Let $\mathcal{C}_s$ (the suffix “s” meaning *source*), $\mathcal{C}_t$ (the suffix “t” meaning *target*) and $\mathcal{B}$ be categories, and let

$$\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t$$

be functors among them (which shall be called a *cospan of categories*). The *comma category*, denoted by $(S \downarrow T)$, is formed by

- the objects of $(S \downarrow T)$ are triples (V, X, h) with $V \in \text{ob}(\mathcal{C}_s)$, $X \in \text{ob}(\mathcal{C}_t)$ and $h \in \text{hom}_{\mathcal{B}}(S(V), T(X))$,
- the morphisms from (V, X, h) to (V', X', h') are pairs (g_s, g_t) where $g_s : V \rightarrow V'$ is a morphism in $\mathcal{C}_s$ and $g_t : X \rightarrow X'$ is a morphism in $\mathcal{C}_t$ such that the following diagram commutes in the category $\mathcal{B}$:

$$\begin{array}{ccc} S(V) & \xrightarrow{S(g_s)} & S(V') \\ h \downarrow & & \downarrow h' \\ T(X) & \xrightarrow{T(g_t)} & T(X'). \end{array}$$

Definition 2.5. Let $\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t$ be a cospan and let $(S \downarrow T)$ be the above comma category associated to the cospan. We define the canonical *projection functors* as follows:

- (1) $\pi_t : (S \downarrow T) \rightarrow \mathcal{C}_t$ is defined by
 - for an object (V, X, h) , $\pi_t((V, X, h)) := X$,
 - for a morphism $(g_s, g_t) : (V, X, h) \rightarrow (V', X', h')$, $\pi_t((g_s, g_t)) := g_t$.
- (2) $\pi_s : (S \downarrow T) \rightarrow \mathcal{C}_s$ is defined by
 - for an object (V, X, h) , $\pi_s((V, X, h)) := V$,
 - for a morphism $(g_s, g_t) : (V, X, h) \rightarrow (V', X', h')$, $\pi_s((g_s, g_t)) := g_s$.

Namely a *cospan of categories* $\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t$ induces a *span of categories* $\mathcal{C}_s \xleftarrow{\pi_s} (S \downarrow T) \xrightarrow{\pi_t} \mathcal{C}_t$.

Definition 2.6. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor of two categories. Then, for an object $B \in \text{ob}(\mathcal{D})$ the *fiber category* of F over B , denoted by $F^{-1}(B)$, is defined to be the category consisting of

- the objects $X \in \text{ob}(\mathcal{C})$ such that $F(X) = B$,
- for such objects X, X' , morphisms $f : X \rightarrow X'$ such that $F(f) = \text{id}_B$.

In other words, more precisely, this category should be denoted by $F^{-1}(B, \text{id}_B)$.

Example 2.7. As above, let us consider a cospan of categories and its associated span of categories:

$$\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t, \quad \mathcal{C}_s \xleftarrow{\pi_s} (S \downarrow T) \xrightarrow{\pi_t} \mathcal{C}_t.$$

- (1) For an object $X \in \mathcal{C}_t$, the fiber category $\pi_t^{-1}(X)$ is nothing but the *S-over category* $(S \downarrow T(X))$, whose objects are *objects S-over T(X)*, i.e., the triple (V, X, h) , and for two triples (V, X, h) and (V', X, h') a morphism from (V, X, h) to (V', X, h') is $g_s \in \text{hom}_{\mathcal{C}_s}(V, V')$ such that the following triangle commutes:

$$\begin{array}{ccc} S(V) & \xrightarrow{S(g_s)} & S(V') \\ & \searrow h & \swarrow h' \\ & T(X) & \end{array}$$

- (2) Furthermore, if $\mathcal{C}_s = \mathcal{B}$ and $S = \text{id}_{\mathcal{B}}$ is the identity functor, then the above *S-over category* $(S \downarrow X)$ is the standard *over category* $(\mathcal{B} \downarrow X)$, whose objects are objects over X , i.e., morphisms $h : V \rightarrow X$, and for two morphisms $h : V \rightarrow X$ and $h' : V' \rightarrow X$ a morphism from $h : V \rightarrow X$ to $h' : V' \rightarrow X$ is

$g \in \text{hom}_{\mathcal{B}}(V, V')$ such that the following triangle commutes:

$$\begin{array}{ccc} V & \xrightarrow{g} & V' \\ & \searrow h \quad \swarrow h' & \\ & X. & \end{array}$$

Remark 2.8. For an object $V \in \mathcal{C}_s$, the fiber category $\pi_s^{-1}(S)$ is nothing but the T -under category $(S(V) \downarrow T)$, whose objects are *objects T -under $S(V)$* , i.e., the triple (V, X, h) , and for two triples (V, X, h) and (V, X', h') a morphism from (V, X, h) to (V, X', h') is $g_t \in \text{hom}_{\mathcal{C}_t}(X, X')$ such that the following triangle commutes:

$$\begin{array}{ccc} & S(V) & \\ h \swarrow & & \searrow h' \\ T(X) & \xrightarrow{T(g_t)} & T(X'). \end{array}$$

Similarly, we can think of the T -under category $(V \downarrow T)$ and the under category $(V \downarrow \mathcal{B})$, but we do not write them down here, since we do not use them below in the rest of the paper.

Proposition 2.9. Let $\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t$ be a cospan of categories. Then a morphism $f \in \text{hom}_{\mathcal{C}_t}(X_1, X_2)$ gives rise to the functor between the corresponding fiber categories:

$$T(f)_* : \pi_t^{-1}(X_1) \rightarrow \pi_t^{-1}(X_2),$$

which is defined by

- (1) For an object (V, X_1, h) , $T(f)_*((V, X_1, h)) := (V, X_2, T(f) \circ h)$.
- (2) For a morphism $(g_s, \text{id}_{X_1}) : (V, X_1, h) \rightarrow (V', X_1, h')$ with $g_s \in \text{hom}_{\mathcal{C}_s}(V, V')$,
 $T(f)_*((g_s, \text{id}_{X_1})) := (g_s, \text{id}_{X_2}) : (V, X_2, T(f) \circ h) \rightarrow (V', X_2, T(f) \circ h')$.

$$\begin{array}{ccccc} S(V) & & \xrightarrow{g} & & S(V') \\ & \searrow h & & \swarrow h' & \\ & T(X_1) & & & \\ & \downarrow T(f) & & & \\ T(f) \circ h & & & & T(f) \circ h' \\ & \searrow & & \swarrow & \\ & T(X_2) & & & \end{array}$$

Lemma 2.10. Let $\mathcal{B}$ be a category with a coproduct $\sqcup$, with the coproduct of any two objects and an initial object chosen, and let $(\mathcal{C}_s, \sqcup)$ be a symmetric monoidal category. Let $\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t$ be a cospan of categories with $S : (\mathcal{C}_s, \sqcup) \rightarrow (\mathcal{B}, \sqcup)$ a strict monoidal functor and $T : \mathcal{C}_t \rightarrow \mathcal{B}$ just a functor. Then for each object $X \in \text{ob}(\mathcal{C}_t)$, the fiber category $\pi_t^{-1}(X)$, i.e. the S -over category $(S \downarrow T(X))$ becomes also a symmetric monoidal category with

$$(V, X, h) \sqcup (V', X, h') := (V \sqcup V', X, h + h').$$

Corollary 2.11. Let the situation be as above. A morphism $f \in \text{hom}_{\mathcal{C}_t}(X_1, X_2)$ gives rise to the canonical group homomorphism

$$T(f)_* : K(\pi_t^{-1}(X_1)) \rightarrow K(\pi_t^{-1}(X_2)),$$

and

$$K(\pi_t^{-1}(-)) : \mathcal{C}_t \rightarrow \mathcal{AB}$$

is a covariant functor from the category $\mathcal{C}_t$ to the category of abelian groups.

After these simple observations, we can introduce the following notions:

Definition 2.12 (*Generalized relative Grothendieck groups with respect to a cospan of categories*).

Let $\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t$ be a cospan of categories as in Lemma 2.10.

- (1) The Grothendieck group of the fiber category of the projection functor $\pi_t : (S \downarrow T) \rightarrow \mathcal{C}_t$ is denoted by

$$K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X)) := K(\pi_t^{-1}(X))$$

and called the *generalized (S, T) -relative Grothendieck group of $X \in \mathcal{C}_t$* .

- (2) If $\mathcal{C}_t = \mathcal{B}$ and $T = id_{\mathcal{B}}$, then $K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$ is simply denoted by $K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/X)$.

- (3) If $S = T = id_{\mathcal{C}_s} : \mathcal{C}_s \rightarrow \mathcal{C}_s$ is the identity functor, then the above $id_{\mathcal{C}}$ -relative Grothendieck group $K(\mathcal{C}_s \xrightarrow{id_{\mathcal{C}_s}} \mathcal{C}_s/X)$ is simply denoted by $K(\mathcal{C}_s/X)$ and called the *relative Grothendieck group of X* or the *relative Grothendieck group of $\mathcal{C}_s$ over X* .

All these relative Grothendieck groups are covariant functors from $\mathcal{C}_t$ to $\mathcal{AB}$.

Remark 2.13. If $T(X) = pt$ is a terminal object in the category $\mathcal{B}$, then all the above relative Grothendieck groups are isomorphic to the Grothendieck group $K(\mathcal{C}_s)$ of the symmetric monoidal category $(\mathcal{C}_s, \sqcup)$:

$$K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X) = pt) \cong K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/pt) \cong K(\mathcal{C}_s/pt) \cong K(\mathcal{C}_s).$$

Proposition 2.14. Let $\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t$ be a cospan of categories as in Lemma 2.10, $\Phi_b : \mathcal{B} \rightarrow \mathcal{B}'$ be a covariant functor preserving (chosen) coproducts and set $S' := \Phi_b \circ S$ and $T' := \Phi_b \circ T$. Then we get the canonical natural transformation from the functor $K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(-)) : \mathcal{C}_t \rightarrow \mathcal{AB}$ to the functor $K(\mathcal{C}_s \xrightarrow{S'} \mathcal{B}'/T'(-)) : \mathcal{C}_t \rightarrow \mathcal{AB}$:

$$\Phi_* : K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(-)) \rightarrow K(\mathcal{C}_s \xrightarrow{S'} \mathcal{B}'/T'(-)),$$

i.e., for a morphism $f \in \text{hom}_{\mathcal{C}_t}(X_1, X_2)$ the following diagram commutes in the category $\mathcal{AB}$:

$$\begin{array}{ccc} K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X_1)) & \xrightarrow{\Phi_*} & K(\mathcal{C}_s \xrightarrow{S'} \mathcal{B}'/T'(X_1)) \\ T(f)_* \downarrow & & \downarrow T'(f)_* \\ K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X_2)) & \xrightarrow{\Phi_*} & K(\mathcal{C}_s \xrightarrow{S'} \mathcal{B}'/T'(X_2)). \end{array}$$

Here $\Phi_* : K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X)) \rightarrow K(\mathcal{C}_s \xrightarrow{S'} \mathcal{B}'/T'(X))$ is defined by

$$\Phi_*([([V, X, h])]) := [(V, X, \Phi_b(h))].$$

Proof. It suffices to show the commutativity $f_*\Phi_* = \Phi_*f_*$ in the square diagram above. First we observe the following morphisms:

$$\begin{array}{c} S'(V) = \Phi_b(S(V)) \\ \downarrow \Phi_b(h) \\ T'(X_1) = \Phi_b(T(X_1)) \\ \downarrow T'(f) = \Phi_b(T(f)) \\ T'(X_2) = \Phi_b(T(X_2)) \end{array}$$

Then we get for $[(V, X_1, h)] \in K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X_1))$:

$$\begin{aligned}
T'(f)_* \Phi_*([(V, X_1, h)]) &= T'(f)_*([(V, X_1, \Phi_b(h))]) \\
&= [(V, X_2, T'(f) \circ \Phi_b(h))] \\
&= [(V, X_2, \Phi_b(T(f)) \circ \Phi_b(h))] \\
&= [(V, X_2, \Phi_b(T(f) \circ h))] \\
&= \Phi_*([(V, X_2, T(f) \circ h)]) \\
&= \Phi_* T(f)_*([(V, X_1, h)]).
\end{aligned}$$

□

As explained in the introduction, when it comes to a categorification of an additive invariant α with values in an additive covariant functor $\mathcal{H}$, we need to consider α -isomorphism class $[(V, X, h)]_\alpha$. Then all the previous results hold even if we replace $[(V, X, h)]$ and $K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(-))$ with $[(V, X, h)]_\alpha$ and $K/\alpha(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(-))$, respectively. And we get the following result:

Theorem 2.15. (A “categorification” of an additive invariant) *Let $\mathcal{B}$ be a category with a coproduct $\sqcup$, with the coproduct of any two objects and an initial object chosen, and let $(\mathcal{C}_s, \sqcup)$ be a symmetric monoidal category. Consider a cospan of categories $\mathcal{C}_s \xrightarrow{S} \mathcal{B} \xleftarrow{T} \mathcal{C}_t$, with $S : (\mathcal{C}_s, \sqcup) \rightarrow (\mathcal{B}, \sqcup)$ a strict monoidal functor and $T : \mathcal{C}_t \rightarrow \mathcal{B}$ just a functor. Let $\mathcal{H} : \mathcal{B} \rightarrow \mathcal{AB}$ be an additive functor on $\mathcal{B}$ with $T' = \mathcal{H} \circ T$. Then an additive invariant α on $Obj(\mathcal{C}_s)$ with values in $\mathcal{H}$ induces a natural transformation on $\mathcal{C}_t$:*

$$\tau_\alpha : K_\alpha(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(-)) \rightarrow T'(-); \tau_\alpha([(V, X, h)]_\alpha) := h_*(\alpha(V)).$$

Assume T is full, and consider one of the cases (i) or (ii) as in Remark 1.2. Then τ_α is the unique natural transformation satisfying

$$\tau_\alpha([V, S(V), \text{id}_{S(V)}]_\alpha) = \alpha(V) \text{ or } \tau_\alpha([V, V, \text{id}_{S(V)}]_\alpha) = \alpha(V).$$

Proof. Note that τ_α is well-defined, since we consider α -isomorphism classes so that it does not depend on the chosen representative of the class $[(V, X, h)]_\alpha$. Then τ_α becomes a group homomorphism by the definition of an additive invariant and the functoriality of $\mathcal{H}$:

$$\begin{aligned}
\tau_\alpha\left([(V, X, h)]_\alpha + [(V', X, h')]_\alpha\right) &= \tau_\alpha([(V, X, h)]_\alpha \sqcup [(V', X, h')]_\alpha) \\
&= \tau_\alpha([(V \sqcup V', X, h + h')]_\alpha) \\
&= (h + h')_*(\alpha(V \sqcup V')) \\
&= (h + h')_*\left((i_{f(V)})_* \alpha(V) + (i_{f(V')})_* \alpha(V')\right) \\
&= (h + h')_*(i_{f(V)})_*(\alpha(V)) + (h + h')_*(i_{f(V')})_*(\alpha(V')) \\
&= h_*(\alpha(V)) + h'_*(\alpha(V')) \\
&= \tau_\alpha([(V, X, h)]_\alpha) + \tau_\alpha([(V', X, h')]_\alpha),
\end{aligned}$$

since $(h + h') \circ i_{f(V)} = h$ and $(h + h') \circ i_{f(V')} = h'$. Finally assume T is full so that there is a morphism $f \in \text{hom}_{\mathcal{C}_t}(S(V), X)$ in the case of (i), or $f \in \text{hom}_{\mathcal{C}_t}(V, X)$ in the case of (ii), with $T(f) = h : S(V) \rightarrow T(X)$. Then

$$f_*([(V, S(V), \text{id}_{S(V)})]_\alpha) = [(V, X, h)]_\alpha \text{ or } f_*([(V, V, \text{id}_{S(V)})]_\alpha) = [(V, X, h)]_\alpha,$$

which implies the uniqueness statement. □

3. A CATEGORIFICATION OF AN ADDITIVE HOMOLOGY CLASS

In the rest of the paper we deal with $\mathcal{B}$ a category of spaces (with some possible extra structures), with the co-product $\sqcup$ given by the usual disjoint union. Examples are the category $\mathcal{TOP}_{(lc)}$ of (locally compact) topological spaces, the category $\mathcal{C}_{(co)}^\infty$ of (closed oriented) C^∞ -manifolds, or the category $\mathcal{V}_k$ of algebraic varieties (i.e. reduced separated schemes of finite type) over a base field k . And the additive covariant functor $\mathcal{H} : \mathcal{B} \rightarrow \mathcal{AB}$ is most of the time a suitable homology theory, like usual homology $H_*(-; R)$ with coefficients in some commutative ring R or a generalized homomolgy H_* , in case $\mathcal{B} = \mathcal{TOP}$. Moreover, for $\mathcal{B} = \mathcal{TOP}_{lc}, \mathcal{C}^\infty$ or $\mathcal{V}_k$, we often consider them as categories only with respect to proper morphisms, with $\mathcal{H}$ the (even degree) Borel-Moore homology $H_{(2)*}^{BM}(-; R)$ with coefficients in some commutative ring R for $\mathcal{B} = \mathcal{TOP}_{lc}, \mathcal{C}^\infty$, or $\mathcal{H} = CH_*$ (resp. $CH_* \otimes R$) the Chow groups (with coefficients in R) for $\mathcal{B} = \mathcal{V}_k$. Similarly, if we further restrict ourselves to projective morphisms in the algebraic context, then $\mathcal{H} : \mathcal{V}_k \rightarrow \mathcal{AB}$ could also be a suitable Borel-Moore functor in the sense of [LM].

The corresponding generalized relative Grothendieck groups only with respect to proper morphisms in $\mathcal{B}$ (and α a corresponding additive invariant) are then denoted by

$$K_{(\alpha)}^{prop}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(-)), K_{(\alpha)}^{prop}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/-) \quad \text{and} \quad K_{(\alpha)}^{prop}(\mathcal{C}_s/-).$$

Then one has a tautological group homomorphism (by just forgetting the properness condition)

$$K^{prop}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X)) \rightarrow K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X)),$$

whose image is the subgroup of the generalized relative Grothendieck group $K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X))$ generated by isomorphism classes $[(V, X, h)]$ with $h : S(V) \rightarrow T(X)$ being a *proper* map. If we assume that $S(V)$ is compact (resp. complete in the algebraic context) for every $V \in ob(\mathcal{C}_s)$, then we have

$$K^{prop}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X)) = K(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(X)).$$

Now our base category $\mathcal{B}$ of spaces is not only a symmetric monoidal category with respect to the disjoint union $\sqcup$ (with unit the initial empty space $\emptyset$), but also with respect to the product of spaces $\times$ (with unit the terminal point space $\{pt\}$, given by $Spec(k)$ in the algebraic context). Moreover, these structures are compatible in the sense that

$$(X \sqcup X') \times Y = (X \times Y) \sqcup (X' \times Y) \quad \text{and} \quad Y \times (X \sqcup X') = (Y \times X) \sqcup (Y \times X'),$$

with $\emptyset \times Y = \emptyset = Y \times \emptyset$. Similarly the class of proper (or projective) morphisms in $\mathcal{B}$ (in the algebraic context) is stable under products $\times$. And we want to discuss the multiplicativity properties of our transformations

$$\tau_\alpha : K_\alpha^{(prop)}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/-) \rightarrow \mathcal{H}$$

associated to an additive invariant on objects in $ob(\mathcal{C}_s)$ with values in a suitable additive functor $\mathcal{H} : \mathcal{B} \rightarrow \mathcal{AB}$ on the category $\mathcal{B}$, which may be functorial only with respect to proper (or projective) morphisms. Here we consider for simplicity only the most important case that $\mathcal{C}_t = \mathcal{B}$ and $T = id_{\mathcal{C}_t}$. Then it is easy to see the following:

Proposition 3.1. (1) *Assume that $\mathcal{C}_s$ is also a symmetric monoidal category with respect to a product $\times$, such that $S : \mathcal{C} = \mathcal{C}_s \rightarrow \mathcal{B}$ is strict monoidal with respect to $\sqcup$ as well as $\times$ (e.g. S is the inclusion of a subcategory stable under $\sqcup$ and $\times$). Then $K^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/X)$ has a functorial bilinear cross product structure:*

$$\times : K^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/X) \times K^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/Y) \rightarrow K^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/X \times Y);$$

$$[(V, X, h)] \times [(W, Y, k)] := [(V \times W, X \times Y, h \times k)],$$

with $\times \circ (f_* \times g_*) = (f \times g)_* \circ \times$ for all (proper or projective) morphisms f, g in $\mathcal{B}$.

(2) *Assume that the additive functor $\mathcal{H} : \mathcal{B} \rightarrow \mathcal{AB}$ is endowed with a bilinear cross product*

$$\boxtimes : \mathcal{H}(X) \times \mathcal{H}(Y) \rightarrow \mathcal{H}(X \times Y)$$

such that $\boxtimes \circ (f_* \times g_*) = (f \times g)_* \circ \boxtimes$ for all (proper or projective) morphisms f, g in $\mathcal{B}$. Consider an additive invariant α on objects in $\text{ob}(\mathcal{C})$ with values in $\mathcal{H}$, which is multiplicative in the sense that

$$\alpha(V \times V') = \alpha(V) \boxtimes \alpha(V') \quad \text{for all } V, V' \in \text{ob}(\mathcal{C}).$$

Then $K_\alpha^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/-)$ also gets a functorial bilinear cross product structure in the same way as before and the associated natural transformation $\tau_\alpha : K_\alpha^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/-) \rightarrow \mathcal{H}$ given in Theorem 2.15 commutes with the cross product, i.e., the following diagram commutes:

$$\begin{array}{ccc} K_\alpha^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/X) \times K_\alpha^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/Y) & \xrightarrow{\times} & K_\alpha^{(prop)}(\mathcal{C} \xrightarrow{S} \mathcal{B}/X \times Y) \\ \tau_\alpha \times \tau_\alpha \downarrow & & \downarrow \tau_\alpha \\ \mathcal{H}(X) \times \mathcal{H}(Y) & \xrightarrow{\boxtimes} & \mathcal{H}(X \times Y). \end{array}$$

Proof. We only need to prove the commutativity of the last diagram, which follows from

$$\begin{aligned} \tau_\alpha([(V, X, h)]_\alpha \times [(W, Y, k)]_\alpha) &= \tau_\alpha([(V \times W, X \times Y, h \times k)]_\alpha) \\ &= (h \times k)_*(\alpha(V \times W)) \\ &= (h \times k)_*(\alpha(V) \boxtimes \alpha(W)) \\ &= h_*(\alpha(V)) \boxtimes k_*(\alpha(W)) \\ &= \tau_\alpha([(V, X, h)]_\alpha) \boxtimes \tau_\alpha([(W, Y, k)]_\alpha). \end{aligned}$$

□

Let us illustrate this result in some examples. First we consider the differential-topological context with $\mathcal{B} = \mathcal{TOP}_{lc}$ the category of locally compact topological spaces, and $\mathcal{C}$ the category $\mathcal{C}_{(o)}^\infty$ or $\mathcal{C}_C^\infty$ of all differentiable (oriented) or stable complex $\mathcal{C}^\infty$ -manifolds, with $S : \mathcal{C} \rightarrow \mathcal{B}$ the forget functor. Of course S commutes with $\sqcup$ and $\times$ for $\mathcal{C} = \mathcal{C}^\infty$ the category of $\mathcal{C}^\infty$ -manifolds. Moreover, any such manifold V has a fundamental class $[V] \in H_*^{BM}(V; \mathbb{Z}_2)$ in Borel-Moore homology with $\mathbb{Z}_2$ -coefficients. And this fundamental class is additive and multiplicative:

$$[V \sqcup V'] = [V] + [V'] \quad \text{and} \quad [V \times V'].$$

When it comes to an oriented (or a stable complex) $\mathcal{C}^\infty$ -manifold V , then this has a fundamental class

$$[V] \in H_*^{BM}(V; \mathbb{Z})$$

in Borel-Moore homology with $\mathbb{Z}$ -coefficients. And we view them as a category with the same morphisms as for the underlying $\mathcal{C}^\infty$ -manifolds, i.e. with $\mathcal{C}^\infty$ -maps between them (so that diffeomorphisms are the isomorphisms). Then the disjoint union $V \sqcup V'$ or product $V \times V'$ of two oriented (or stable complex) $\mathcal{C}^\infty$ -manifolds V, V' can also be oriented (or given the structure of a stable complex) $\mathcal{C}^\infty$ -manifold. And there is a natural choice for this so that the fundamental class $[-]$ becomes additive and multiplicative as before. In this way we also get the symmetric monoidal structures $\sqcup$ and $\times$ on the category of oriented (or a stable complex) $\mathcal{C}^\infty$ -manifolds, with the forget functor S to $\mathcal{TOP}_{lc}$ (or also to $\mathcal{C}^\infty$) commuting with these structures. Also note that

$$s_*[V \times V'] = (-1)^{\dim(V) \cdot \dim(V')} [V' \times V] \in H_{\dim(V) + \dim(V')}^{BM}(V' \times V; \mathbb{Z})$$

for two *connected* oriented manifolds V, V' and $s : V \times V' \xrightarrow{\sim} V' \times V$ the symmetry isomorphism, since $H_*^{BM}(-; \mathbb{Z})$ is *graded-commutative* with respect to the usual cross product $\boxtimes$.

Corollary 3.2. *Let $\mathcal{C}$ be the category $\mathcal{C}_{(o)}^\infty$ or $\mathcal{C}_C^\infty$ of all (oriented) or stable complex smooth manifolds. Consider a contravariant functorial characteristic class*

$$cl(E) \in H^*(-; \mathbb{Z}_2) \quad \text{or} \quad cl(E) \in H^{2*}(-; R)$$

of (isomorphism classes of) real (oriented) or complex vector bundles, which is multiplicative and normalized, i.e.:

$$cl(E \oplus F) = cl(E) \cup cl(F) \quad \text{and} \quad 1 = cl^0(E) \in H^0(-; R).$$

For a smooth (oriented or stable complex) manifold V , let

$$\alpha(V) := \mathcal{cl}(TV) \cap [V] \in H_*^{BM}(-; R).$$

Then the invariant α is additive and multiplicative. By Theorem 2.15, there exists a unique natural transformation

$$\tau_{\mathcal{cl}} : K_{\alpha}^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{TOP}_{lc}/-) \rightarrow H_*^{BM}(-; R)$$

such that for a smooth (oriented or stable complex) manifold V

$$\tau_{\mathcal{cl}}([(V, V, id_V)]_{\alpha}) = \mathcal{cl}(TV) \cap [V].$$

And $\tau_{\mathcal{cl}}$ is also multiplicative (by Proposition 3.1), i.e.:

$$\tau_{\mathcal{cl}}([(V, X, h)]_{\alpha} \times [(W, Y, k)]_{\alpha}) = \tau_{\mathcal{cl}}([(V, X, h)]_{\alpha}) \boxtimes \tau_{\mathcal{cl}}([(W, Y, k)]_{\alpha}).$$

Proof. The invariant α is additive by the functoriality of $\mathcal{cl}$ and the projection formula for the inclusions $i : V \rightarrow V \sqcup V'$ and $i' : V' \rightarrow V \sqcup V'$:

$$\begin{aligned} \alpha(V \sqcup V') &= \mathcal{cl}(T(V \sqcup V')) \cap [V \sqcup V'] \\ &= \mathcal{cl}(T(V \sqcup V')) \cap (i_*[V] + i'_*[V']) \\ &= i_*(i^*\mathcal{cl}(T(V \sqcup V')) \cap [V]) + i'_*(i'^*\mathcal{cl}(T(V \sqcup V')) \cap [V']) \\ &= i_*(\mathcal{cl}(TV) \cap [V]) + i'_*(\mathcal{cl}(TV') \cap [V']) \\ &= i_*\alpha(V) + i'_*\alpha(V'). \end{aligned}$$

The invariant α is multiplicative by the functoriality and multiplicativity of $\mathcal{cl}$:

$$\begin{aligned} \alpha(V \times V') &= \mathcal{cl}(T(V \times V')) \cap [V \times V'] \\ &= (\mathcal{cl}(TV) \boxtimes \mathcal{cl}(TV')) \cap ([V] \boxtimes [V']) \\ &= (\mathcal{cl}(TV) \cap [V]) \boxtimes (\mathcal{cl}(TV') \cap [V']) \\ &= \alpha(V) \boxtimes \alpha(V'). \end{aligned}$$

Note that in the third equality there is no sign appearing, since we only consider even degree characteristic classes $\mathcal{cl}$ in the context of oriented or stable complex manifolds. $\square$

Remark 3.3. In the context of oriented manifolds in the above corollary, (V, X, h) and (V', X, h) are α -isomorphic if and only if the isomorphism $\phi : V \rightarrow V'$ (with $h = h' \circ S(\phi)$) is orientation preserving. Indeed, if $\phi : V \rightarrow V'$ is orientation preserving, then $S(\phi)_*[V] = [V']$ and $\phi^*TV' \simeq TV$ as oriented vector bundles. Hence we have

$$\begin{aligned} S(\phi)_*(\alpha(V)) &= S(\phi)_*(\mathcal{cl}(TV) \cap [V]) \\ &= S(\phi)_*(S(\phi)^*\mathcal{cl}(TV') \cap [V]) \\ &= \mathcal{cl}(TV') \cap S(\phi)_*[V] \quad (\text{by the projection formula}) \\ &= \mathcal{cl}(TV') \cap [V'] = \alpha(V'). \end{aligned}$$

Hence (V, X, h) and (V', X, h) are α -isomorphic. Conversely, if (V, X, h) and (V', X, h) are α -isomorphic, then we have $S(\phi)_*(\alpha(V)) = \alpha(V')$, i.e., $S(\phi)_*(\mathcal{cl}(TV) \cap [V]) = \mathcal{cl}(TV') \cap [V']$. Since $\mathcal{cl}$ is normalized, we have $\mathcal{cl} = 1 + \dots$, hence $\mathcal{cl}(TV) \cap [V] = [V] + \text{lower dimensional classes}$. Therefore we have

$$S(\phi)_*[V] = [V'].$$

Hence $\phi : V \rightarrow V'$ is orientation preserving.

Similarly α is an isomorphism invariant in the context of unoriented manifolds. In the context of stable complex manifolds, α is at least invariant under a diffeomorphism $\phi : V \rightarrow V'$ of stable complex manifolds who preserves the stable almost complex structure (and therefore also the orientation) in the sense that $\phi^*TV' \simeq TV$ as stable complex vector bundles.

If we consider in Corollary 3.2 all *compact* (oriented) or stable complex smooth manifolds, then we get similar results for any *generalized homology theory* $\mathcal{H}_*$, which has a corresponding fundamental class $[V] \in \mathcal{H}_*(V)$ for a compact (oriented) or stable complex smooth manifold V , e.g. for a complex oriented (co)homology theory and V a stable complex smooth manifold.

Let us now switch to some counterparts in the algebraic geometric context, with $\mathcal{B} = \mathcal{V}_k$ the category of algebraic varieties (i.e. reduced separated schemes of finite type) over a base field k , and $S : \mathcal{C} \rightarrow \mathcal{V}_k$ the inclusion functor of a (full) subcategory $\mathcal{C}$ stable under isomorphisms, disjoint union $\sqcup$ and product $\times$, with $\emptyset, \text{Spec}(k) \in \text{ob}(\mathcal{C})$. First we consider the subcategory $\mathcal{C} = \mathcal{V}_k^{\text{sm}}$ of smooth varieties. The proof of the following result is identical to that of Corollary 3.2.

Corollary 3.4. *Consider the cospan of categories*

$$\mathcal{V}_k^{\text{sm}} \xrightarrow{S} \mathcal{V}_k \xleftarrow{id_{\mathcal{V}_k}} \mathcal{V}_k,$$

together with a contravariant functorial characteristic class

$$cl(E) \in CH^*(-) \otimes R \quad \text{or} \quad cl(E) \in H^{2*}(-; R) \quad \text{for } k = \mathbb{C}$$

of (isomorphism classes of) algebraic vector bundles, which is multiplicative in the sense that

$$cl(E) = cl(E') \cup cl(E'')$$

for any short exact sequence $0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$ of such vector bundles. Here $CH^*(-)$ is the operational Chow cohomology group of [Fu]. For a smooth algebraic manifold V , let

$$\alpha(V) := cl(TV) \cap [V] \in \mathcal{H}_*(-) \otimes R,$$

with $[V]$ the fundamental class of V for $\mathcal{H}_* = CH_*$ the Chow group or H_{2*}^{BM} the even degree Borel-Moore homology in the case of $k = \mathbb{C}$. Then the isomorphism invariant α is additive and multiplicative. By Theorem 2.15, there exists a unique natural transformation

$$\tau_{cl} : K^{\text{prop}}(\mathcal{V}_k^{\text{sm}} \xrightarrow{S} \mathcal{V}_k / -) \rightarrow \mathcal{H}_*(-) \otimes R$$

such that for a smooth algebraic manifold V

$$\tau_{cl}([(V, V, id_V)]) = cl(TV) \cap [V].$$

And τ_{cl} is also multiplicative (by Proposition 3.1), i.e.:

$$\tau_{cl}([(V, X, h)]) \times [(W, Y, k)] = \tau_{cl}([(V, X, h)]) \boxtimes \tau_{cl}([(W, Y, k)]).$$

Remark 3.5. If we only consider projective morphisms and (pure dimensional) quasi-projective smooth varieties, then a similar result holds for $\mathcal{H}_*$ an *oriented Borel-Moore weak homology theory* in the sense of [LM] and cl a multiplicative characteristic class as in [LM, §4.1.8], which is, for a line bundle L , given by a normalized power series in the first Chern class operator of L with respect to $\mathcal{H}_*$:

$$cl(L) = f(\tilde{c}_1(L)), \quad \text{with } f(t) \in 1 + t \cdot \mathcal{H}_*(pt)[[t]].$$

Here the fundamental class of a quasi-projective smooth variety V of pure dimension d is defined as

$$[V] := k^* 1_{pt} \in \mathcal{H}_d(V) \quad \text{for } k : V \rightarrow pt \text{ the constant smooth morphism.}$$

Next we consider the subcategory $\mathcal{C} = \mathcal{V}_k^{(l)pd}$ of (locally) pure-dimensional algebraic varieties over k . Let us first recall the universal property of the *Nash blow-up* $\nu : \hat{X} \rightarrow X$ of a pure d -dimensional algebraic variety $X \in \text{Obj}(\mathcal{V}_k^{pd})$, with $\widehat{TX}$ the *tautological Nash tangent bundle* over $\hat{X}$: Let $\pi : \overline{X} \rightarrow X$ be a proper birational map with a surjection $\pi^* \Omega_X^1 \rightarrow \overline{\Omega}$ to a locally free sheaf $\overline{\Omega}$ of rank d on $\overline{X}$. Then the Nash blow-up $\nu : \hat{X} \rightarrow X$ is universal in the sense that $\pi : \overline{X} \rightarrow X$ factors through $\pi' : \overline{X} \rightarrow \hat{X}$, with $\overline{\Omega} \simeq \pi'^* \hat{\Omega}$, where the tautological Nash tangent bundle $\widehat{TX}$ over $\hat{X}$ corresponds to the dual of $\hat{\Omega}$.

Definition 3.6. Let $c\ell$ be a functorial characteristic class of algebraic vector bundles as in Corollary 3.4. For a pure d -dimensional algebraic variety $X \in \text{Obj}(\mathcal{V}_k^{pd})$, the $c\ell$ -Mather homology class $c\ell_*^{Ma}(X) \in \mathcal{H}_*(X) \otimes R$ is defined to be

$$c\ell_*^{Ma}(X) := \nu_*(c\ell(\widehat{T\overline{X}}) \cap [\widehat{X}]) = \pi_*(c\ell(\overline{T\overline{X}}) \cap [\overline{X}]).$$

Here $\pi : \overline{X} \rightarrow X$ is any proper birational map with a surjection $\pi^*\Omega_X^1 \rightarrow \overline{\Omega}$ to a locally free sheaf $\overline{\Omega}$ of rank d on $\overline{X}$, with the vector bundle $\overline{T\overline{X}}$ corresponding to the dual of $\overline{\Omega}$. This definition is extended to a locally pure-dimensional variety $X \in \text{Obj}(\mathcal{V}_k^{lpd})$ by additivity over the connected components of X .

Note that the second equality in the definition above follows from the projection formula by $\overline{T\overline{X}} \simeq \pi'^*\widehat{T\overline{X}}$ and $\pi'_*[\overline{X}] = [\widehat{X}]$, since $\pi' : \overline{X} \rightarrow \widehat{X}$ is a proper birational map (see [Fu, Example 4.2.9(b)] in the case of the Chern-Mather class c^{Ma} corresponding to $c\ell = c$ the Chern class).

Corollary 3.7. Consider the cospan of categories

$$\mathcal{V}_k^{(l)pd} \xrightarrow{S} \mathcal{V}_k \xleftarrow{id_{\mathcal{V}_k}} \mathcal{V}_k,$$

together with a contravariant functorial characteristic class

$$c\ell(E) \in CH^*(-) \otimes R \quad \text{or} \quad c\ell(E) \in H^{2*}(-; R) \quad \text{for } k = \mathbb{C}$$

of (isomorphism classes of) algebraic vector bundles, which is multiplicative. Here $CH^*(-)$ is the operational Chow cohomology group [Fu, Definition 17.3]. For a (locally) pure-dimensional algebraic variety V , let

$$\alpha(V) := c\ell_*^{Ma}(V) \in \mathcal{H}_*(V) \otimes R,$$

for $\mathcal{H}_* = CH_*$ the Chow group or H_{2*}^{BM} the even degree Borel-Moore homology in the case of $k = \mathbb{C}$.

- (1) The isomorphism invariant α is additive and multiplicative. By Theorem 2.15, there exists a unique natural transformation

$$\tau_{c\ell_*^{Ma}} : K^{prop}(\mathcal{V}_k^{(l)pd} \xrightarrow{S} \mathcal{V}_k / -) \rightarrow \mathcal{H}_*(-) \otimes R$$

such that for a (locally) pure-dimensional algebraic variety V

$$\tau_{c\ell_*^{Ma}}([(V, V, id_V)]) = c\ell_*^{Ma}(V).$$

And $\tau_{c\ell_*^{Ma}}$ is also multiplicative (by Proposition 3.1), i.e.:

$$\tau_{c\ell_*^{Ma}}([(V, X, h)] \times [(W, Y, k)]) = \tau_{c\ell_*^{Ma}}([(V, X, h)]) \boxtimes \tau_{c\ell_*^{Ma}}([(W, Y, k)]).$$

- (2) When $c\ell = c$ is the Chern class, then the following diagram commutes for k of characteristic zero:

$$\begin{array}{ccc} K^{prop}(\mathcal{V}_k^{(l)pd} \xrightarrow{S} \mathcal{V}_k / X) & \xrightarrow{\mathcal{E}u} & F(X) \\ & \searrow \tau_{c_*^{Ma}} & \swarrow c_*^{Ma} \\ & \mathcal{H}_*(X) & \end{array}$$

Here the multiplicative natural transformation $\mathcal{E}u : K^{prop}(\mathcal{V}_k^{(l)pd} \xrightarrow{C} \mathcal{V}_k / X) \rightarrow F(X)$ to the group of constructible functions is defined by the isomorphism invariant $\alpha(V) := Eu_V$, where Eu_V is the local Euler obstruction of the (locally) pure-dimensional variety V . This invariant is also additive and multiplicative.

Proof. (1) That the isomorphism invariant $\alpha(V) := c\ell_*^{Ma}(V)$ is additive follows from the fact, that it commutes with restriction to open subsets (e.g. a connected component). For the multiplicativity we can then assume that V, V' are pure dimensional. Let $\pi : \overline{V} \rightarrow V$ be a proper birational map with a surjection $\pi^*\Omega_X^1 \rightarrow \overline{\Omega}$ to a locally free sheaf $\overline{\Omega}$, and similarly for V' . Then $\pi \times \pi' : \overline{V} \times \overline{V'} \rightarrow V \times V'$ is a proper birational map with a surjection

$$(\pi \times \pi')^*\Omega_{V \times V'}^1 \simeq \pi^*\Omega_V^1 \boxtimes \pi'^*\Omega_{V'}^1 \rightarrow \overline{\Omega} \boxtimes \overline{\Omega'}$$

so that

$$\begin{aligned}
 c\ell_*^{Ma}(V \times V') &= (\pi \times \pi')_* (c\ell(\overline{TV} \boxtimes \overline{TV'}) \cap [\overline{V} \times \overline{V'}]) \\
 &= (\pi \times \pi')_* ((c\ell(\overline{TV}) \cap [\overline{V}]) \boxtimes (c\ell(\overline{TV'}) \cap [\overline{V'}])) \\
 &= (\pi_*(c\ell(\overline{TV}) \cap [\overline{V}])) \boxtimes (\pi'_*(c\ell(\overline{TV'}) \cap [\overline{V'}])) \\
 &= c\ell_*^{Ma}(V) \boxtimes c\ell_*^{Ma}(V').
 \end{aligned}$$

(2) follows from the construction [Mac1] of the Chern class transformation

$$c_*^{Mac} : F(X) \rightarrow H_{2*}^{BM}(X; \mathbb{Z})$$

for the case $k = \mathbb{C}$. MacPherson defined $c_*^{Mac}(Eu_V) = (i_V)_* c_*^{Ma}(V)$ for $i_V : V \rightarrow X$ the inclusion of a pure-dimensional subvariety, with Eu_V his famous local Euler obstruction. Namely, the “constructible function” counterpart of the Chern–Mather homology class has to be the local Euler obstruction, which is one of his key observations. This is extended to locally pure-dimensional subvarieties by additivity over connected components, e.g. the local Euler obstruction is then by definition the sum of the local Euler obstructions of all connected components. The algebraic counterpart of the MacPherson Chern class transformation

$$c_*^{Mac} : F(X) \rightarrow CH_*(X)$$

for a base field k of characteristic zero was constructed in [Ke] (at least if X is embeddable into a smooth variety. The general case can be reduced to this using the method of Chow envelopes as in [Fu, Chapter 18.3]). Moreover, Kennedy also explained in [Ke] that $Eu_V \in F(V)$ is a constructible function, where he used the algebraic definition of the local Euler obstruction as in [Fu, Example 4.2.9] (due to Gonzalez-Sprinberg and Verdier):

$$Eu_V(p) = \int_{\pi^{-1}(p)} c(\overline{TV}|_{\pi^{-1}(p)}) \cap s(\pi^{-1}(p), \overline{V}),$$

with $\pi : \overline{V} \rightarrow V$ and $\overline{TV}$ are as in the definition of the $c\ell$ -Mather homology classes. Here $s(\pi^{-1}(p), \overline{V})$ is the Segre class of the fiber $\pi^{-1}(p)$ in $\overline{V}$ in the sense of [Fu, Chapter 4.2]. Then the multiplicativity of Eu_V follows as in (1) using [Fu, Example 4.2.5]:

$$s(\pi^{-1}(p) \times \pi'^{-1}(p'), \overline{V} \times \overline{V'}) = s(\pi^{-1}(p), \overline{V}) \boxtimes s(\pi'^{-1}(p'), \overline{V'}).$$

Here the bilinear (functorial) cross product

$$\boxtimes : F(X) \times F(Y) \rightarrow F(X \times Y)$$

is just defined by $\beta \boxtimes \beta'((p, p')) := \beta(p) \cdot \beta'(p')$. □

Remark 3.8. Assume that the base field k is of characteristic zero.

- (1) Using resolution of singularities one can show that for a given algebraic variety X there are finitely many irreducible subvarieties V ’s and integers a_V ’s such that

$$\mathbb{1}_X = \sum_{V \subset X} a_V Eu_V, \text{ thus } c_*^{Mac}(\mathbb{1}_X) = \sum_{V \subset X} a_V c_*^{Ma}(V).$$

Whether X is singular or not, $c_*^{Mac}(X) := c_*^{Mac}(\mathbb{1}_X)$ is called MacPherson’s Chern class or Chern–Schwarz–MacPherson class of X (see [BrSc, Ke, Mac1, Schw1, Schw2]). For X complete, it follows from the naturality of the transformation c_*^{Mac} with respect to the proper constant map $X \rightarrow pt$, that the degree of the 0-dimensional component of $c_0^{Mac}(X)$ is equal to the Euler–Poincaré characteristic:

$$\int_X c_0^{Mac}(X) = \chi(X).$$

- (2) Similarly, the degree of the 0-dimensional component of the Chern–Mather class $c_*^{Ma}(X)$ for X pure-dimensional and complete is the Euler–Poincaré characteristic of X weighted by the local Euler obstruction Eu_X :

$$\int_X c_0^{Ma}(X) = \chi(X; Eu_X).$$

For X a connected *complex affine* algebraic variety of pure dimension, the *global Euler obstruction* $Eu(X)$ introduced and studied in [STV] is a suitable “localization” of the 0-dimensional component of the Chern–Mather class $c_*^{Ma}(X)$:

$$H_0(X; \mathbb{Z}) \simeq \mathbb{Z} \ni Eu(X) \mapsto c_0^{Ma}(X) \in H_0^{BM}(X; \mathbb{Z})$$

under the natural map $H_0(X; \mathbb{Z}) \rightarrow H_0^{BM}(X; \mathbb{Z})$.

- (3) The above “ $c\ell$ -Mather class” transformation $\tau_{c\ell^{Ma}} : K^{prop}(\mathcal{V}_k^{(l)pd} \xrightarrow{S} \mathcal{V}_k/-) \rightarrow \mathcal{H}_*(-) \otimes R$ could be considered as a very naïve theory of characteristic classes of possibly singular algebraic varieties (in any characteristic).

So far we dealt with the covariance and multiplicativity of the functor $K_{(\alpha)}(\mathcal{C}_s \xrightarrow{S} \mathcal{B}/T(-))$. Next we discuss the contravariance with respect to (suitable) “smooth morphisms”, where we start with the algebraic geometric context with $\mathcal{B} = \mathcal{V}_k$ the category of algebraic varieties over k and $S : \mathcal{C} \rightarrow \mathcal{V}_k$ the inclusion functor of the (full) subcategories $\mathcal{C} = \mathcal{V}_k^{sm}$ resp. $\mathcal{V}_k^{lci}$ of *smooth* resp. *local complete intersection* varieties.

Here X is called a local complete intersection, if it has a regular closed embedding $i : X \rightarrow M$ into a smooth variety M (i.e. the constant morphism $X \rightarrow pt$ is a local complete intersection morphism in the sense of [Fu, Chapter 6.6]). Then X has an intrinsic virtual tangent bundle

$$TX := i^*TM - N_X M \in K^0(X),$$

with $N_X M$ the normal bundle of the regular embedding $i : X \rightarrow M$, i.e. $TX \in K^0(X)$ doesn’t depend on the choice of this embedding (compare [Fu, Appendix B.7.6]). Of course any smooth variety M is local complete intersection, with TM (the class of) the usual tangent bundle TM (just choose $i = id_M : M \rightarrow M$). Note that the (virtual) tangent bundle commutes with restriction to open subsets (e.g. connected components). Similarly, if $f : X \rightarrow Y$ is a smooth morphism with Y smooth (resp. a local complete intersection), then also X is smooth (resp. a local complete intersection), and in the smooth context we have a short exact sequence of vector bundles

$$(3.9) \quad 0 \rightarrow T_f \rightarrow TX \rightarrow f^*TY \rightarrow 0.$$

with T_f the bundle of tangents to the fiber of the smooth morphism f . In particular

$$(3.10) \quad TX = f^*TY + T_f \in K^0(X),$$

and this equality in the Grothendieck group $K^0(-)$ of algebraic vector bundles even holds for a smooth morphism $f : X \rightarrow Y$ between local complete intersections with TX resp. TY the corresponding virtual tangent bundle (compare e.g. with [FL, Proposition 7.1] as well as [Fu, Appendix B.7]). Finally, the class of smooth morphisms is stable under base-change, with $T_{f'} \simeq h'^*T_f$ for a fiber square as in the following

Lemma 3.11. *The functor $K^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{V}_k/-)$, with $\mathcal{C} = \mathcal{V}_k^{sm}$ (resp. $\mathcal{V}_k^{lci}$) the subcategory of smooth (resp. local complete intersection) varieties, becomes a contravariant functor for smooth morphisms on the category $\mathcal{V}_k$, where for a smooth morphism $f : X \rightarrow Y$ the pullback homomorphism*

$$f^* : K^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{V}_k/Y) \rightarrow K^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{V}_k/X)$$

is defined by

$$f^*([(V, Y, h)]) := [V', X, h'],$$

using the fiber square

$$\begin{array}{ccc} V' & \xrightarrow{f'} & V \\ h' \downarrow & & \downarrow h \\ X & \xrightarrow{f} & Y. \end{array}$$

Of course here we also use the fact that taking such fiber squares commutes with disjoint unions in V resp. V' .

Theorem 3.12 (Verdier-type Riemann–Roch). *Consider the cospan of categories*

$$\mathcal{C} \xrightarrow{S} \mathcal{V}_k \xleftarrow{id_{\mathcal{V}_k}} \mathcal{V}_k,$$

with $\mathcal{C} = \mathcal{V}_k^{sm}$ (resp. $\mathcal{V}_k^{lci}$) the subcategory of smooth (resp. local complete intersection) varieties. Let cl be a contravariant functorial characteristic class

$$cl(E) \in CH^*(-) \otimes R \quad \text{or} \quad cl(E) \in H^{2*}(-; R) \quad \text{for } k = \mathbb{C}$$

of (isomorphism classes of) algebraic vector bundles, which is multiplicative (and normalized in case $\mathcal{C} = \mathcal{V}_k^{lci}$, so that it can also be defined on virtual vector bundles in $K^0(-)$). For a smooth (resp. local complete intersection) variety V , let

$$\alpha(V) := cl(TV) \cap [V] \in \mathcal{H}_*(V) \otimes R$$

be the corresponding (virtual) characteristic homology class, with $\mathcal{H}_* = CH_*$ the Chow group or H_{2*}^{BM} the even degree Borel-Moore homology in case $k = \mathbb{C}$.

The isomorphism invariant α is additive, so that by Theorem 2.15 there exists a unique natural transformation

$$\tau_{cl} : K^{prop}(\mathcal{C} \xrightarrow{C} \mathcal{V}_k / -) \rightarrow \mathcal{H}_*(-) \otimes R,$$

such that for a smooth (resp. local complete intersection) algebraic variety V

$$\tau_{cl}([(V, V, id_V)]) = cl(TV) \cap [V].$$

Then this natural transformation τ_{cl} satisfies the following Verdier-type Riemann–Roch formula: For a smooth morphism $f : X \rightarrow Y$ the following diagram commutes:

$$\begin{array}{ccc} K^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{V}_k / Y) & \xrightarrow{\tau_{cl}} & \mathcal{H}_*(Y) \otimes R \\ f^* \downarrow & & \downarrow cl(T_f) \cap f^* \\ K^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{V}_k / X) & \xrightarrow{\tau_{cl}} & \mathcal{H}_*(X) \otimes R. \end{array}$$

Proof. On one hand we have:

$$\begin{aligned} \tau_{cl}(f^*([(V, Y, h)])) &= \tau_{cl}([(V', X, h')]) \\ &= h'_*(cl(TV') \cap [V']). \end{aligned}$$

On the other hand we have

$$\begin{aligned} cl(T_f) \cap f^*(\tau_{cl}([(V, Y, h)])) \\ = cl(T_f) \cap f^*(h_*(cl(TV) \cap [V])) \end{aligned}$$

For a fiber square

$$\begin{array}{ccc} V' & \xrightarrow{f'} & V \\ h' \downarrow & & \downarrow h \\ X & \xrightarrow{f} & Y \end{array}$$

with $h : V \rightarrow Y$ proper and $f : X \rightarrow Y$ smooth, we have the base change identity (see [Fu, Proposition 1.7]):

$$f^* h_* = h'_* f'^* : \mathcal{H}_*(V) \otimes R \rightarrow \mathcal{H}_*(X) \otimes R.$$

Hence the above equality continues as follows:

$$\begin{aligned} &= c\ell(T_f) \cap h'_* f'^* (c\ell(TV) \cap [V]) \\ &= h'_* (h'^* c\ell(T_f) \cap f'^* (c\ell(TV) \cap [V])) \text{ (by the projection formula)} \\ &= h'_* (c\ell(T_{f'}) \cap f'^* (c\ell(TV) \cap [V])) \text{ (by } T_{f'} \simeq h'^* T_f) \\ &= h'_* (c\ell(T_{f'}) \cap (c\ell(f'^* TV) \cap f'^* [V])) \text{ (by functoriality of } c\ell(-) \cap) \\ &= h'_* ((c\ell(T_{f'}) \cup c\ell(f'^* TV)) \cap [V']) \\ &= h'_* (c\ell(TV') \cap [V']) \end{aligned}$$

by the multiplicativity of $c\ell$ and (3.9) resp. (3.10). Of course we also used the relation $[V'] = f'^*[V]$ for the fundamental classes. Therefore we get that

$$\tau_{c\ell}(f^*([(V, Y, h)])) = c\ell(T_f) \cap f^*(\tau_{c\ell}([(V, Y, h)])),$$

and the above diagram of the theorem commutes. $\square$

Remark 3.13. If we consider only projective morphisms and (pure dimensional) quasi-projective local complete intersection (resp. smooth) varieties, then a similar Verdier-type Riemann–Roch formula holds for $\mathcal{H}_*$ an *oriented Borel-Moore (weak) homology theory* in the sense of [LM] and $c\ell$ a multiplicative characteristic class as in [LM, §7.4.1 resp. §4.1.8], which is, for a line bundle L , given by a normalized power series in the first Chern class operator of L with respect to $\mathcal{H}_*$:

$$c\ell(L) = f(\tilde{c}_1(L)), \quad \text{with } f(t) \in 1 + t \cdot \mathcal{H}_*(pt)[[t]].$$

Here the fundamental class of a quasi-projective local complete intersection (resp. smooth) variety V of pure dimension d is defined as

$$[V] := k^* 1_{pt} \in \mathcal{H}_d(V) \quad \text{for } k : V \rightarrow pt \text{ the constant local complete intersection (resp. smooth) morphism}$$

so that $[V'] = f'^*[V]$ for a smooth morphism $f' : V' \rightarrow V$ by functoriality of f'^* .

Remark 3.14. Assume that the base field k is of characteristic zero.

(1) The interesting thing about the motivic Hirzebruch class or MacPherson Chern class transformation

$$T_{y*} : K^{prop}(\mathcal{V}_k^{sm} \xrightarrow{S} \mathcal{V}_k / -) \rightarrow \mathcal{H}_*(-) \otimes \mathbb{Q}[y] \quad \text{or} \quad c_*^{Mac} : K^{prop}(\mathcal{V}_k^{sm} \xrightarrow{S} \mathcal{V}_k / -) \rightarrow \mathcal{H}_*(-)$$

is for example, that in the above discussions, the naïve relative Grothendieck group $K^{prop}(\mathcal{V}_k^{sm} \xrightarrow{S} \mathcal{V}_k / X)$ can be replaced by the much smaller and more interesting relative Grothendieck group $K_0(\mathcal{V}_k / X)$, by imposing one more *additivity relation*, as recalled in the introduction:

$$[V \xrightarrow{h} X] = [W \xrightarrow{h|_W} X] + [V \setminus W \xrightarrow{h|_{V \setminus W}} X],$$

with $W \subset V$ is a closed subvariety of V . So these two transformations factorize over the tautological surjective transformation

$$K^{prop}(\mathcal{V}_k^{sm} \xrightarrow{S} \mathcal{V}_k / -) \rightarrow K_0(\mathcal{V}_k / -).$$

In particular the multiplicativity result of Corollary 3.4 and the Verdier-type Riemann–Roch formula of Theorem 3.12 are true for the motivic Hirzebruch class and the MacPherson Chern class transformation (compare [BSY]).

- (2) The corresponding Hirzebruch class $T_{y*}(X) = T_{y*}(id_X)$ and MacPherson Chern class $c_*^{Mac}(X) = c_*^{Mac}(id_X)$ of an algebraic variety X is not only invariant under an isomorphism, but also under a proper (geometric) bijection:

$$f_*(T_{y*}(X)) = T_{y*}(Y) \quad \text{and} \quad f_*(c_*^{Mac}(X)) = c_*^{Mac}(Y)$$

for a proper morphism $f : X \rightarrow Y$ such that the induced map $f : X(\bar{k}) \rightarrow Y(\bar{k})$ is a bijection of sets (with $\bar{k}$ an algebraic closure of k), since then $f_*([id_X]) = [id_Y]$ by Noetherian induction using additivity and generic smoothness of f . Of course for MacPherson's Chern class $c_*^{Mac}(X) = c_*^{Mac}(\mathbb{1}_X)$ this also follows from $f_*\mathbb{1}_X = \mathbb{1}_Y \in F(Y)$ for a proper (geometric) bijection $f : X \rightarrow Y$. Similarly, Baum–Fulton–MacPherson's Todd class $td_*^{BFM}(X) = td_*^{BFM}([\mathcal{O}_X])$ is invariant under a such a proper (geometric) bijection (for k of any characteristic), since $f_*[\mathcal{O}_X] = [\mathcal{O}_Y] \in G_0(Y)$. Finally, Goresky–MacPherson's or Cappell–Shaneson's homology L -class $L_*(X) = L_*^{CS}([\mathcal{IC}_X])$ of a (locally) pure-dimensional compact complex algebraic variety X is not only invariant under a proper bijection, but more generally under taking the normalization of X .

Let us finish this section with a counterpart in the differential-topological context, with $\mathcal{B} = \mathcal{C}^\infty$ the category of $\mathcal{C}^\infty$ -manifolds and $S : \mathcal{C} \rightarrow \mathcal{C}^\infty$ the forget functor from the category $\mathcal{C} = \mathcal{C}_{(o)}^\infty$ or $\mathcal{C}_{\mathbb{C}}^\infty$ of all differentiable (oriented) or stable complex $\mathcal{C}^\infty$ -manifolds.

Lemma 3.15. *The functor $K^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{C}^\infty/-)$ for $\mathcal{C} = \mathcal{C}_{(o)}^\infty$ or $\mathcal{C}_{\mathbb{C}}^\infty$ becomes a contravariant functor for a ((complex) oriented) submersion $f : X \rightarrow Y$ of $\mathcal{C}^\infty$ -manifolds. Here the pullback homomorphism*

$$f^* : K^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{C}^\infty/Y) \rightarrow K^{prop}(\mathcal{C} \xrightarrow{C} \mathcal{C}^\infty/X)$$

is defined by

$$f^*([(V, Y, h)]) := [V', X, h'],$$

using the fiber square

$$\begin{array}{ccc} V' & \xrightarrow{f'} & V \\ h' \downarrow & & \downarrow h \\ X & \xrightarrow{f} & Y, \end{array}$$

with TV' oriented (or given a stable complex structure) by the short exact sequence

$$0 \rightarrow T_{f'} \rightarrow TV' \rightarrow f'^*TV \rightarrow 0,$$

if TV and the bundle of tangents to the fibers T_f (and therefore also $T_{f'} \simeq h'^*T_f$) are oriented (or stable complex).

Then the proof of the following result is identical to that in the algebraic geometric context:

Theorem 3.16 (Verdier-type Riemann–Roch for smooth manifolds). *Consider the cospan of categories*

$$\mathcal{C} \xrightarrow{S} \mathcal{C}^\infty \xleftarrow{id_{\mathcal{C}^\infty}} \mathcal{C}^\infty,$$

with $\mathcal{C} = \mathcal{C}_{(o)}^\infty$ or $\mathcal{C}_{\mathbb{C}}^\infty$ the category of all differentiable (oriented) or stable complex $\mathcal{C}^\infty$ -manifolds. Let $c\ell$ be a contravariant functorial multiplicative and normalized characteristic class

$$c\ell(E) \in H^*(-; \mathbb{Z}_2) \quad \text{or} \quad c\ell(E) \in H^{2*}(-; R)$$

of (isomorphism classes of) real (oriented) or complex vector bundles. For a smooth (oriented or stable complex) manifold V , let

$$\alpha(V) := c\ell(TV) \cap [V] \in H_*^{BM}(V; R).$$

The invariant α is additive, so that by Theorem 2.15 there exists a unique natural transformation

$$\tau_{c\ell} : K_\alpha^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{C}^\infty/-) \rightarrow H_*^{BM}(-; R),$$

such that for a smooth (oriented or stable complex) manifold V

$$\tau_{c\ell}([(V, V, id_V)]) = c\ell(TV) \cap [V].$$

Then this natural transformation $\tau_{c\ell}$ satisfies the following Verdier-type Riemann–Roch formula: For a ((complex) oriented) submersion $f : X \rightarrow Y$ of $\mathcal{C}^\infty$ -manifolds, the following diagram commutes:

$$\begin{array}{ccc} K_\alpha^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{C}^\infty/Y) & \xrightarrow{\tau_{c\ell}} & H_*^{BM}(Y; R) \\ f^* \downarrow & & \downarrow c\ell(T_f) \cap f^* \\ K_\alpha^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{C}^\infty/X) & \xrightarrow{\tau_{c\ell}} & H_*^{BM}(X; R). \end{array}$$

4. EXAMPLES

In this last section we discuss some results, questions and problems related to some very specific examples, which also fit with our natural transformations associated to some additive (homology) invariants.

4.1. The case of the fundamental class. Let us first consider taking the fundamental class $[-] \in H_*^{BM}(-; R)$ for $R = \mathbb{Z}_2$ (or $R = \mathbb{Z}$) on the category $\mathcal{C}_{(o)}^\infty$ of (oriented) C^∞ -manifolds. The fundamental class $\alpha := [-]$ is certainly an additive and multiplicative homology class and by Corollary 3.2 we have a unique natural transformation

$$\tau_{[-]} : K_\alpha^{prop}(\mathcal{C} \xrightarrow{S} \mathcal{TOP}_{lc}/-) \rightarrow H_*^{BM}(-; R)$$

such that $\tau_{[-]}([(V, V, id_V)]) = [V]$ for a smooth (oriented) manifold V . Here an α -isomorphism $\phi : V \rightarrow V'$ is just an (orientation preserving) diffeomorphism by Remark 3.3.

Of course such an additive (and multiplicative) fundamental homology class $\alpha := [-]$ is also available on bigger categories of spaces, like the categories $\mathcal{C}_{(o)}^0$ of (oriented) topological manifolds, or $\mathcal{C}_{(o)}^{pseudo}$ the category of (oriented) triangulated or topological pseudo-manifolds, whose morphisms are by definition just continuous maps of the underlying topological spaces (so that an isomorphism is nothing but a homeomorphism). If we restrict ourselves to compact spaces (as classically often done), then these fundamental classes live in usual homology $H_*(-; R)$ for $R = \mathbb{Z}_2$ (or $R = \mathbb{Z}$ in the oriented context). Moreover such an additive and multiplicative fundamental class $\alpha = [-] \in H_*(-; R)$ is also available for the category $\mathcal{C}_{(o)}^{Poincaré}$ of finite (oriented) Poincaré complexes, i.e., topological spaces V (like homology-manifolds) which satisfy Poincaré duality

$$\cap[V] : H^*(V; R) \xrightarrow{\sim} H_*(V; R)$$

for a suitable fundamental class $[V] \in H_*(V; R)$. So we can turn these categories $\mathcal{C}$ as before into symmetric monoidal categories with respect to disjoint unions $\sqcup$ (and also products $\times$) such that the forgetful functor $f : \mathcal{C} \rightarrow \mathcal{TOP}_{lc}$ is strictly monoidal. Moreover, an α -isomorphism $\phi : V \rightarrow V'$ in this context is just an (orientation preserving) homeomorphism.

Then the classical *Steenrod's realization problem* can be reinterpreted as the problem of asking for the surjectivity of the group homomorphism $\tau_{[-]} : K_\alpha(\mathcal{C} \xrightarrow{f} \mathcal{TOP}/X) \rightarrow H_*(X; R)$ for a topological space X . Here the following results are well known (see [Rud2, Sul2]).

Theorem 4.1. (1) ([Th1] and [Rud1, Chapter IV, Theorem 7.33])

$$\tau_{[-]} : K(\mathcal{C}_c^\infty \xrightarrow{f} \mathcal{TOP}/X) \rightarrow H_*(X; \mathbb{Z}_2)$$

is surjective, i.e. every $\mathbb{Z}_2$ -homology class can be realized by a compact smooth manifold.

(2) ([Th1] and [Rud1, Chapter IV, Theorem 7.37]) The following composed map is surjective:

$$\tau_{[-]} : K_\alpha(\mathcal{C}_{co}^\infty \xrightarrow{f} \mathcal{TOP}/X) \rightarrow H_*(X; \mathbb{Z}) \xrightarrow{proj.} \bigoplus_{0 \leq i \leq 6} H_i(X; \mathbb{Z}),$$

i.e. every $\mathbb{Z}$ -homology class in degree $i \leq 6$ can be realized by a compact oriented smooth manifold.

(3) ([Lev]) *The following composed map is surjective:*

$$\tau_{[-]} : K_\alpha(\mathcal{C}_{co}^{Poincaré} \xrightarrow{f} \mathcal{TOP}/X) \rightarrow H_*(X; \mathbb{Z}) \xrightarrow{proj.} \bigoplus_{i \neq 3} H_i(X; \mathbb{Z}),$$

i.e. every $\mathbb{Z}$ -homology class in degree $i \neq 3$ can be realized by a finite oriented Poincaré complex.

(4) ([Sul1] and [Rud1, Chapter VIII, Example 1.25(a)])

$$\tau_{[-]} : K_\alpha(\mathcal{C}_{co}^{pseudo} \xrightarrow{f} \mathcal{TOP}/X) \rightarrow H_*(X; \mathbb{Z})$$

is surjective, i.e. every $\mathbb{Z}$ -homology class can be realized by a compact oriented pseudo-manifold.

4.2. The case of the Stiefel–Whitney class. Let V be a differentiable manifold and let $c\ell_*(V) \in H_*^{BM}(V; \mathbb{Z}_2)$ be the Poincaré dual $c\ell(TV) \cap [V]$ of a multiplicative and normalized functorial characteristic class $c\ell(E) \in H^*(-, \mathbb{Z}_2)$ of (isomorphism classes of) real vector bundles. $\alpha(V) := c\ell_*(V)$ is clearly an additive (and multiplicative) homology class, and we have by Corollary 3.2 a unique natural transformation

$$\tau_{c\ell_*} : K^{prop}(\mathcal{C}^\infty \xrightarrow{S} \mathcal{TOP}_{lc}/-) \rightarrow H_*^{BM}(-; \mathbb{Z}_2)$$

such that $\tau_{c\ell_*}([(V, V, id_V)]) = c\ell_*(V)$ for a smooth manifold V . In particular the Stiefel–Whitney class $c\ell = w$ is a typical one.

If we restrict ourselves to the category $\mathcal{V}_{\mathbb{R}}$ of real algebraic varieties and let $\mathcal{V}_{\mathbb{R}}^{sm}$ be its full subcategory of smooth real algebraic varieties, then we have a more geometric “realization” of the natural transformation w_* on the category $\mathcal{V}_{\mathbb{R}}$ through $\mathbb{Z}_2$ -valued semi-algebraic constructible functions:

$$\begin{array}{ccc} K^{prop}(\mathcal{V}_{\mathbb{R}}^{sm} \xrightarrow{t} \mathcal{V}_{\mathbb{R}}/X) & \xrightarrow{const} & F(X; \mathbb{Z}_2) \\ & \searrow w_* & \swarrow w_* \\ & H_*^{BM}(X, \mathbb{Z}_2) & \end{array}$$

Here the multiplicative natural transformation $const$ is defined by the isomorphism invariant $\alpha(V) := 1_V$, which is additive and multiplicative. The Stiefel–Whitney class transformation $w_* : F(X; \mathbb{Z}_2) \rightarrow H_*^{BM}(X, \mathbb{Z}_2)$ was first constructed by Sullivan [Sul1] in the pl-context. For a new approach in the semi-algebraic (or subanalytic) context through the theory of “conormal or characteristic cycles” see [FMc].

Remark 4.2. For a compact topological manifold V , Thom constructed a Whitney class using a relation with Steenrod squares [Th2] (see [Mac2]). Let us denote this Thom–Whitney class in homology by $w_*^{Th}(V) \in H_*(V; \mathbb{Z}_2)$, which for a compact smooth manifold V agrees with the Stiefel–Whitney class $w_*(V)$ above. Then also $\alpha(V) := w_*^{Th}(V)$ is an additive (and multiplicative) invariant, so that we have a natural transformation

$$\tau_{w_*}^{Th} : K(\mathcal{C}_c^0 \xrightarrow{f} \mathcal{TOP}/-) \rightarrow H_*(-; \mathbb{Z}_2)$$

defined by

$$\tau_{w_*}^{Th}([(V, X, h)]) = h_* w_*^{Th}(V).$$

If we consider the above Whitney class transformation

$$w_* : K(\mathcal{C}_c^\infty \xrightarrow{f} \mathcal{TOP}/-) \rightarrow H_*(-; \mathbb{Z}_2)$$

for compact smooth C^∞ -manifolds, then this is surjective by Theorem 4.1(i), since $w_*(V) = [V] + \dots$ equals the fundamental class plus lower order terms. So a natural problem is to find for a given compact topological manifold X a class $\beta \in K(\mathcal{C}_c^\infty \xrightarrow{f} \mathcal{TOP}/X)$ such that

$$w_*(\beta) = w_*^{Th}(X).$$

4.3. The case of the Pontryagin and L-class. Let V be an oriented differentiable manifold and let $\mathcal{cl}_*(V) \in H_{2*}^{BM}(V; \mathbb{Q})$ be the Poincaré dual $\mathcal{cl}(TV) \cap [V]$ of a multiplicative and normalized functorial characteristic class $\mathcal{cl}(E) \in H^{2*}(-, \mathbb{Q})$ of (isomorphism classes of) oriented real vector bundles. $\alpha(V) := \mathcal{cl}_*(V)$ is clearly an additive (and multiplicative) homology class, and by Corollary 3.2 we have a unique natural transformation

$$\tau_{\mathcal{cl}_*} : K_{\alpha}^{prop}(\mathcal{C}_o^{\infty} \xrightarrow{S} \mathcal{TOP}_{lc}/-) \rightarrow H_{2*}^{BM}(-; \mathbb{Q})$$

such that $\tau_{\mathcal{cl}_*}([(V, V, id_V)]) = \mathcal{cl}_*(V)$ for a smooth oriented manifold V . In particular the Pontryagin class $\mathcal{cl} = p$ and the Hirzebruch-Thom L -class $\mathcal{cl} = L$ are typical ones.

If we restrict ourselves to the category $\mathcal{V}_{\mathbb{C},c}$ of compact complex algebraic varieties and let $\mathcal{V}_{\mathbb{C},c}^{sm}$ be its full subcategory of compact smooth complex algebraic varieties, then we have a more geometric “realization” of the natural transformation L_* on the category $\mathcal{V}_{\mathbb{C},c}$ through *Cappell–Shaneson–Youssin’s cobordism groups* $\Omega_*(X)$ (see [CS], [You]) and the motivic relative Grothendieck group as mentioned in the introduction (compare with [BSY] for more details):

$$\begin{array}{ccc} K(\mathcal{V}_{\mathbb{C},c}^{sm} \xrightarrow{L} \mathcal{V}_{\mathbb{C},c}/X) & \xrightarrow{taut} & K_0(\mathcal{V}_{\mathbb{C}}/X) \\ L_* \downarrow & & \downarrow sd \\ H_{2*}(X, \mathbb{Q}) & \xleftarrow{L_*^{CS}} & \Omega(X). \end{array}$$

Here the (composed) multiplicative natural transformation

$$sd : K(\mathcal{V}_{\mathbb{C},c}^{sm} \xrightarrow{L} \mathcal{V}_{\mathbb{C},c}/X) \rightarrow \Omega(X)$$

is defined by the isomorphism invariant $\alpha(V) := [\mathbb{Q}_V[dim V]] \in \Omega(V)$, which is additive and multiplicative.

Remark 4.3. For a compact triangulated $\mathbb{Q}$ -homology manifold V , Thom constructed in [Th3] an L -class in $H^{2*}(V; \mathbb{Q}) \simeq H_{2*}(V; \mathbb{Q})$ using a relation with the signature (see [Mac2, Sul2]), so that for a compact oriented smooth manifold it agrees with the usual L -class $L(TV)$ of the tangent bundle by the famous *Hirzebruch signature theorem*. Through the development of *Intersection (co)homology*, this approach was further extended (by e.g. Goresky-MacPherson [GM] and Cappell-Shaneson [CS]) to more general singular spaces like compact (locally) pure-dimensional complex algebraic varieties. But then one has to view this class as a homology L -class in $H_{2*}(V; \mathbb{Q})$. Nevertheless, in Thom’s original approach for a compact triangulated $\mathbb{Q}$ -homology manifold V only, one can use Poincaré duality to view this as a cohomology class in $H^{2*}(V; \mathbb{Q})$. Using the cup-product structure on cohomology, he was then able to define also a Pontryagin class $p(V) \in H^{2*}(V; \mathbb{Q})$ so that for a compact oriented smooth manifold it agrees with the usual Pontryagin class $p(TV)$ of the tangent bundle.

Let us call the corresponding homology classes the Thom–Pontryagin and Thom L -class, denoted by

$$p_*^{Th}(X), L_*^{Th}(X) \in H_{2*}(X; \mathbb{Q}).$$

If we consider the above Pontryagin- or L -class transformation for $\mathcal{cl} = p$ or L in the context of compact oriented smooth manifolds

$$\tau_{\mathcal{cl}_*} : K_{\alpha}(\mathcal{C}_{co}^{\infty} \xrightarrow{S} \mathcal{TOP}/-) \rightarrow H_{2*}(-; \mathbb{Q}),$$

then for a given compact triangulated $\mathbb{Q}$ -homology manifold X it is a very interesting problem to find a class $\beta \in K(\mathcal{C}_{co}^{\infty} \xrightarrow{S} \mathcal{TOP}/X)$ such that

$$p_*(\beta) = p_*^{Th}(X) \quad \text{or} \quad L_*(\beta) = L_*^{Th}(X).$$

Note that the rationalized group homomorphism $\tau_{\mathcal{cl}_*} \otimes \mathbb{Q}$ is surjective by [Th1] (compare [Rud1, Chapter IV, Theorem 7.36]), since $\mathcal{cl}_*(V) = [V] + \dots$ equals the fundamental class plus lower order terms.

4.4. The case of the Chern class. Let V be a stable complex differentiable manifold and let $c\ell_*(V) \in H_{2*}^{BM}(V, R)$ be the Poincaré dual $c\ell(TV) \cap [V]$ of a multiplicative and normalized functorial characteristic class $c\ell(E) \in H^{2*}(-, R)$ of (isomorphism classes of) complex vector bundles. $\alpha(V) := c\ell_*(V)$ is clearly an additive (and multiplicative) homology class, and we have by Corollary 3.2 a unique natural transformation

$$\tau_{c\ell_*} : K^{prop}(\mathcal{C}_{\mathbb{C}}^{\infty} \xrightarrow{S} \mathcal{TOP}_{lc}/-) \rightarrow H_{2*}^{BM}(-; R)$$

such that $\tau_{c\ell_*}([(V, V, id_V)]) = c\ell_*(V)$ for a smooth stable complex manifold V . In particular the Chern class $c\ell = c$ is a typical one.

If we restrict ourselves to the category $\mathcal{V}_{\mathbb{C}}$ of complex algebraic varieties and let $\mathcal{V}_{\mathbb{C}}^{sm}$ be its full subcategory of smooth complex algebraic varieties, then we have a more geometric “realization” of the natural transformation c_* on the category $\mathcal{V}_{\mathbb{C}}$ through constructible functions and the motivic relative Grothendieck group as mentioned in the introduction (see [BSY] for more details):

$$\begin{array}{ccc} K^{prop}(\mathcal{V}_{\mathbb{C}}^{sm} \xrightarrow{L} \mathcal{V}_{\mathbb{C}}/X) & \xrightarrow{taut} & K_0(\mathcal{V}_{\mathbb{C}}/X) \\ c_* \downarrow & & \downarrow const \\ H_{2*}^{BM}(X, \mathbb{Z}) & \xleftarrow{c_*^{Mac}} & F(X). \end{array}$$

Here the (composed) multiplicative natural transformation

$$const : K^{prop}(\mathcal{V}_{\mathbb{C}}^{sm} \xrightarrow{L} \mathcal{V}_{\mathbb{C}}/X) \rightarrow F(X)$$

is defined by the isomorphism invariant $\alpha(V) := \mathbb{1}_V \in F(V)$, which is additive and multiplicative.

4.5. The case of Chern classes of other types. Let $\mathcal{V}_{\mathbb{C}}^{emb}$ be the subcategory of (complex) algebraic varieties embeddable into smooth varieties and let $c_*^{FJ}(X)$ resp., $c_*^F(X)$ be Fulton–Johnson’s Chern class resp., Fulton’s canonical class defined for such an embeddable (complex) algebraic variety: $c_*^{FJ}(X)$ ([Fu, Example 4.2.6 (c)]) is defined by

$$c_*^{FJ}(X) := c(TM|_X) \cap s(\mathcal{N}_X M),$$

where TM is the tangent bundle of M and $s(\mathcal{N}_X M)$ is the Segre class of the conormal sheaf $\mathcal{N}_X M$ of X in M [Fu, §4.2]. Fulton’s canonical class $c_*^F(X)$ ([Fu, Example 4.2.6 (a)]) is defined by

$$c_*^F(X) := c(TM|_X) \cap s(X, M),$$

where $s(X, M)$ is the relative Segre class [Fu, §4.2]. For a local complete intersection variety X we also have a normal bundle $N_X M$ in M , from which we can define the virtual tangent bundle T_X of X (as in §3) by

$$T_X := TM|_X - N_X M \in K^0(X).$$

As shown in [Fu, Example 4.2.6], for a local complete intersection variety X in a non-singular variety M , these two Chern classes are both equal to the virtual Chern class

$$c_*^{FJ}(X) = c_*^F(X) = c(T_X) \cap [X].$$

Moreover, both isomorphism invariants $\alpha(V) := c_*^{FJ}(V)$ and $c_*^F(V)$ are additive for $V \in ob(\mathcal{V}_{\mathbb{C}}^{emb})$, so that there exists unique natural transformations on the category $\mathcal{V}_{\mathbb{C}}$:

$$\tau_{c_*^F}, \tau_{c_*^{FJ}} : K^{prop}(\mathcal{V}_{\mathbb{C}}^{emb} \xrightarrow{L} \mathcal{V}_{\mathbb{C}}/-) \rightarrow H_{2*}^{BM}(-; \mathbb{Z}),$$

such that

$$\tau_{c_*^F}([V \xrightarrow{id_V} V]) = c_*^F(V) \quad \text{resp.} \quad \tau_{c_*^{FJ}}([V \xrightarrow{id_V} V]) = c_*^{FJ}(V)$$

for $V \in ob(\mathcal{V}_{\mathbb{C}}^{emb})$. Finally, using Chow groups $CH_*(X)$ as a corresponding homology theory, all of this remains true over any base field k (instead of working over $\mathbb{C}$ with Borel-Moore homology), with the invariant $\alpha(V) = c_*^F(V)$ for $V \in ob(\mathcal{V}_k^{emb})$ also multiplicative (as follows from [Fu, Example 4.2.5]).

Acknowledgements. We would like to thank Paolo Aluffi and Markus Banagl for useful discussions, and also Yuli B. Rudyak for providing the precise reference for Theorem 4.1. Finally we would like to thank Lê Dũng Tráng for useful suggestions and the referee for his/her careful reading and valuable suggestions and comments.

REFERENCES

- [Al] P. Aluffi, *Singular schemes of hypersurfaces*, Duke J. Math. **180** (1995), 325 – 351.
- [BFM] P. Baum, W. Fulton and R. MacPherson, *Riemann–Roch for singular varieties*, Publ. Math. I.H.E.S. **45** (1975), 101–145.
- [BLSS] J.-P. Brasselet, D. Lehman, J. Seade and T. Suwa, *Milnor classes of local complete intersection*, Tran. Amer. Math. Soc. **354** (2001), 1351–1371.
- [BSY] J.-P. Brasselet, J. Schürmann and S. Yokura, *Hirzebruch classes and motivic Chern classes for singular spaces*, Journal of Topology and Analysis, Vol. **2**, No.1 (2010), 1–55.
- [BrSc] J.-P. Brasselet and M.-H. Schwartz, *Sur les classes de Chern d’une ensemble analytique complexe*, Astérisque **82–83**(1981), 93–148.
- [CMSS] S. E. Cappell, L. Maxim, J. L. Shaneson and J. Schürmann, *Hirzebruch classes of hypersurfaces*, Adv. in Math. **225** (2010), 2616–2647.
- [CS] S. E. Cappell and J. L. Shaneson, *Stratifiable maps and topological invariants*, J. Amer. Math. Soc. **4** (1991), 521–551.
- [FMc] J. Fu and C. McCrory, *Stiefel–Whitney classes and the conormal cycle of a singular variety*, Trans. Amer. Math. Soc. **349** (1997), 809–835.
- [Fu] W. Fulton, *Intersection Theory*, Springer Verlag, 1981.
- [FJ] W. Fulton and K. Johnson, *Canonical classes on singular varieties*, Manus. Math. **32** (1980), 381–389.
- [FL] W. Fulton and S. Lang, *Riemann–Roc Algebra*, Springer Verlag New York, 1985.
- [FM] W. Fulton and R. MacPherson, *Categorical frameworks for the study of singular spaces*, Memoirs of Amer. Math. Soc. **243**, 1981.
- [GM] M. Goresky and R. MacPherson, *Intersection homology theory*, Topology, **149** (1980), 155–162.
- [Hir] F. Hirzebruch, *Topological Methods in Algebraic Geometry*, 3rd ed. (1st German ed. 1956), Springer-Verlag, 1966.
- [HBJ] F. Hirzebruch, T. Berger and R. Jung, *Manifolds and Modular Forms*, Vieweg, 1992.
- [Ke] G. Kennedy, *MacPherson’s Chern classes of singular varieties*, Com. Algebra. **9** (1990), 2821–2839.
- [LM] M. Levine and F. Morel, *Algebraic Cobordism*, Springer–Verlag Berlin Heidelberg, 2007.
- [Lev] N. Levitt, *Applications of Poincaré duality spaces to the topology of manifolds*, Topology **11** (1972), 205–221.
- [MacL] S. MacLane, *Categories for the Working Mathematician*, Springer–Verlag New York, 1998.
- [Mac1] R. MacPherson, *Chern classes for singular algebraic varieties*, Ann. of Math. **100** (1974), 423–432.
- [Mac2] R. MacPherson, *Characteristic classes for singular varieties*, Proceedings of the 9th Brazilian Mathematical Colloquium (Poços de Caldas, 1973), Vol II, Instituto de Matematica Pura e Aplicada, São Paulo (1977), 321–327.
- [PP] A. Parusiński and P. Pragacz, *Characteristic classes of hypersurfaces and characteristic cycles*, J. Algebraic Geometry, **10** (2001), 63–79.
- [Rud1] Y. B. Rudyak, *On Thom Spectra, Orientability, and Cobordism*, Springer Monographs in Mathematics, Springer, 1998.
- [Rud2] Yu. B. Rudyak, *Steenrod problem*, Springer Online Reference Works, Encyclopaedia of Mathematics (Edited by Michiel Hazewinkel), <http://eom.springer.de/S/s087540.htm>
- [Sch] J. Schürmann, *Characteristic classes of mixed Hodge modules*, in “Topology of Stratified Spaces”, MSRI Publications **58**, Cambridge Univ. Press (2010), 419–470,
- [SY] J. Schürmann and S. Yokura, *A survey of characteristic classes of singular spaces*, in “Singularity Theory” (Denis Cheniot et al, ed.), Dedicated to Jean-Paul Brasselet on his 60th birthday, Proceedings of the 2005 Marseille Singularity School and Conference, World Scientific (2007), 865–952,
- [Schw1] M.-H. Schwartz, *Classes caractéristiques définies par une stratiification d’une variété analytique complexe*, C. R. Acad. Sci. Paris **t. 260** (1965), 3262–3264, 3535–3537.
- [Schw2] M.-H. Schwartz, *Classes et caractères de Chern des espaces linéaires*, C. R. Acad. Sci. Paris Sér. I. Math. **295** (1982), 399–402.
- [STV] J. Seade, T. Mihai and A. Verjovsky, *Global Euler obstruction and polar invariants*, Math. Ann. **333** (2005), no.2, 393–403.
- [Sul1] D. Sullivan, *Geometric periodicity and the invariants of manifolds*, Manifolds–Amsterdam 1970 (Proc. Nuffic Summer School), Springer Lecture Notes in Mathematics, Vol. **197** (1971), 44–75.
- [Sul2] D. Sullivan, *René Thom’s work on geometric homology and bordism*, Bulletin Amer. Math. Soc. (New Series), **41**, No. 3 (2004), 341–350.
- [Th1] R. Thom, *Quelques propriétés globales des variétés différentiables*, Comm. Math. Helv., **28** (1954), 17–86.
- [Th2] R. Thom, *Espaces fibrés en sphères et carrés de Steenrod*, Ann. Sci. Ecole Norm. Sup., **69** (1952), 109–182.
- [Th3] R. Thom, *Les classes caractéristique de Pontrjagin des variétés triangulés*, Symposium Internacional de Topologia Algebraica, UN-ESCO (1958), 54–67.
- [Yo1] S. Yokura, *Polar classes and Segre classes on singular projective varieties*, Trans. Amer. Math. Soc., 298 (1986), 169–191,
- [Yo2] S. Yokura, *On Cappell-Shaneson’s homology L-class of singular algebraic varieties*, Trans. Amer. Math. Soc. **347** (1995), 1005–1012
- [Yo3] S. Yokura, *On characteristic classes of complete intersections*, “Algebraic Geometry - Hirzebruch 70” (ed. by Pragacz, Szurek and Wisniewski) Contemporary Mathematics Amer. Math. Soc., **241** (1999), 349–369
- [Yo4] S. Yokura, *Motivic characteristic classes*, in “Topology of Stratified Spaces”, MSRI Publications **58**, Cambridge Univ. Press (2010), 375–418,

- [Yo5] S. Yokura, *Motivic Milnor classes*, Journal of Singularities, **1** (2010), 39–59.
- [Yo6] S. Yokura, *Natural transformations associated to additive homology classes*, Kyoto University RIMS Kôkyûroku **1764** (2011), 30–47.
- [You] B. Youssin, *Witt Groups of Derived Categories*, K-Theory, **11** (1997), 373–395.

JÖRG SCHÜRMANN: WESTF. WILHELMS-UNIVERSITÄT, MATHEMATISCHES INSTITUT, EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY

E-mail address: jschuerm@math.uni-muenster.de

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, FACULTY OF SCIENCE, KAGOSHIMA UNIVERSITY, 21-35 KORIMOTO 1-CHOME, KAGOSHIMA 890-0065, JAPAN

E-mail address: yokura@sci.kagoshima-u.ac.jp